

ANTACID MINT- aluminum hydroxide, magnesium hydroxide, demethicone suspension

Walgreens

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active Ingredients

Aluminum hydroxide (equiv. to dried gel, USP) 200 mg
Magnesium hydroxide 200 mg
Simethicone 20 mg

Purpose

Antacid

Antacid

Antigas

Uses

- For the relief of
 - acid indigestion
 - heartburn
 - sour stomach
 - upset stomach associated with these symptoms
 - pressure and bloating commonly referred to as gas

Warnings

Do not take

more than 16 teaspoonfuls in a 24-hour period or use the maximum dosage for more than 2 weeks except under the advice and supervision of a physician

Ask a doctor before use if you have

- kidney disease
- a magnesium restricted diet

Ask a doctor or pharmacist before use if you are

presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if

symptoms last for more than 2 weeks

If pregnant or breast-feeding

ask a health professional before use.

Keep out of the reach of children.

In case of overdose, get medical help or contact a Poison Controls Center (1-800-222-1222) right away.

Directions

- shake well before using
- do not take more than 16 teaspoonfuls in 24 hours or use the maximum dosage for more than 2 week
- adults and children 12 years of age and older: take 2 to 4 teaspoonfuls four times a day or as directed by a physician
- children under 12 years of age: do not use

Other Information

- each teaspoon (5 mL) contains: magnesium 80 mg, potassium 5 mg, and sodium 2 mg
- store between 20-25°C (68-77°F). Avoid freezing

Inactive Ingredients

benzyl alcohol, carboxymethylcellulose sodium, flavors, glycerin, microcrystalline cellulose, potassium citrate, propylene glycol, purified water, sorbitol, saccharin sodium xanthan gum

Principal Display Panel

Compare to Maalox® Advanced Regular Strength Active ingredient^{††}

Antacid +

Anti-Gas Liquid

ALUMINUM HYDROXIDE 200 mg / ANTACID

MAGNESIUM HYDROXIDE 200 mg / ANTACID

SIMETHICONE 20 mg /ANTI-GAS

REGULAR STRENGTH

- Fast relief of heartburn, pressure, bloating & acid indigestion

FL OZ (mL)

MINT FLAVOR

††This product is not manufactured or distributed by Novartis Consumer Health, Inc., distributor of Maalox® Advanced Regular Strength.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

**DISTRIBUTED BY: WALGREENS CO.,
200 WILMOT RD., DEERFIELD, IL 60015**

walgreens.com

Package Label

The image shows the packaging for Walgreens Antacid + Anti-Gas Liquid. The top part of the packaging is blue with the Walgreens logo in red. Below the logo, it says "Compare to Maalox® Advanced Regular Strength Active Ingredients!" and "NDC 0363-0432-26". The main text on the packaging reads "Antacid + Anti-Gas Liquid" in large white letters. Below that, it lists the active ingredients: "ALUMINUM HYDROXIDE 200 mg / ANTACID", "MAGNESIUM HYDROXIDE 200 mg / ANTACID", and "SIMETHICONE 20mg / ANTI-GAS". A blue banner says "REGULAR STRENGTH". Below that, it lists benefits: "Fast relief of heartburn, pressure, bloating & acid indigestion" and "26 FL OZ (769 mL)". There is an image of green leaves and the text "MINT FLAVOR". On the right side of the packaging, there is a vertical label "PLD-A457B LB005702".

Drug Facts	
Active ingredients (in each 5 mL = 1 teaspoon)	Purposes
Aluminum hydroxide (equiv. to dried gel USP) 200 mg.....	Antacid
Magnesium hydroxide 200 mg.....	Antacid
Simethicone 20 mg.....	Antigas

Uses ■ for the relief of
■ acid indigestion ■ heartburn ■ sour stomach
■ upset stomach associated with these symptoms
■ pressure and bloating commonly referred to as gas

Warnings
Do not take more than 16 teaspoonfuls in a 24-hour period or use the maximum dosage for more than 2 weeks except under the advice and supervision of a physician.

Ask a doctor before use if you have
■ kidney disease ■ a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ shake well before using
■ do not take more than 16 teaspoonfuls in 24 hours or use the maximum dosage for more than 2 weeks

Drug Facts (continued)
■ adults and children 12 years of age and older: take 2 to 4 teaspoonfuls four times a day or as directed by a physician
■ children under 12 years of age: do not use

Other information
■ each teaspoon (5 mL) contains: magnesium 80 mg, potassium 5 mg, and sodium 2 mg
■ store between 20-25°C (68-77°F). Avoid freezing.

Inactive ingredients benzyl alcohol, carboxymethylcellulose sodium, flavor, glycerin, microcrystalline cellulose, potassium citrate, propylene glycol, purified water, saccharin sodium, sorbitol, xanthan gum

Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

Walgreens Pharmacist Recommended
Walgreens Pharmacist Survey
††This product is not manufactured or distributed by Novartis Consumer Health, Inc., distributor of Maalox® Advanced Regular Strength.
DISTRIBUTED BY: WALGREEN CO., 200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2018 Walgreen Co.
ITEM 272286 W10023-0418-F

3 11917 20012 5
PLD-A457B LB005703 ORG0418-F2

WALGREENS Antacid and Anti-Gas

ANTACID MINT

aluminum hydroxide, magnesium hydroxide, demithicone suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0432
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	200 mg in 5 mL
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM HYDROXIDE	200 mg in 5 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	20 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
BENZYL ALCOHOL (UNII: LKG8494WBH)	
GLYCERIN (UNII: PDC6A3C0OX)	
POTASSIUM CITRATE (UNII: EE90ONI6FF)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SACCHARIN SODIUM ANHYDROUS (UNII: I4807BK602)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0432-26	769 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/31/2018	12/29/2023
2	NDC:0363-0432-12	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/31/2018	12/29/2023

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part332	03/31/2018	12/29/2023

Labeler - Walgreens (008965063)

Revised: 4/2022

Walgreens