

**LOW DOSE ASPIRIN- aspirin tablet, chewable**  
**Walgreen Company**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Walgreens 44-397**

**Active ingredient (in each tablet)**

Aspirin 81 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever

**Uses**

for the temporary relief of minor aches and pains or as recommended by your doctor

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- have 3 or more alcoholic drinks every day while using this product
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- take more or for a longer time than directed

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease

- you have asthma
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- new symptoms occur
- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts more than 10 days
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

***Directions***

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

***Other information***

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

***Inactive ingredients***

cherry flavor, corn starch, D&C red #27 aluminum lake, D&C red #30 aluminum lake, dextrans, sodium saccharin

**Principal Display Panel**

**VALUE PACK (3 BOTTLES) x 36 = 108**

Well at

**Walgreens**

WALGREENS PHARMACIST RECOMMENDED<sup>□</sup>

NDC 0363-0397-07

**CHEWABLE • LOW DOSE**

**Aspirin 81**

81 mg / Pain Reliever (NSAID)

• For aspirin regimen users

**HEART HEALTH**

**108**

CHEWABLE

TABLETS

(3 BOTTLES: 36 TABLETS PER BOTTLE)

**CHEERRY FLAVOR**

**Compare to Bayer<sup>®</sup> Chewable Low Dose Aspirin active ingredient<sup>□□</sup>**

<sup>□</sup>Walgreens Pharmacist Survey Study, November 2010.

<sup>□□</sup>This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Bayer<sup>®</sup> Chewable Low Dose Aspirin.

50844 REV0613A39707

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS  
BROKEN OR MISSING**

**DISTRIBUTED BY: WALGREEN CO.**

**200 WILMOT RD., DEERFIELD, IL 60015**

**100% SATISFACTION GUARANTEED**

walgreens.com

©2012 Walgreen Co.



VALUE PACK X 36 = 108

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

**Drug Facts**

**Active ingredient (in each tablet)**

Aspirin 81 mg (NSAID) ..... Pain reliever  
 \*nonsteroidal anti-inflammatory drug

**Purpose**

**Uses** for the temporary relief of minor aches and pains or as recommended by your doctor

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

B-2201-397-07-TBR  
REV0613A39707

**Drug Facts (continued)**

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives ■ facial swelling
- shock ■ asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- have 3 or more alcoholic drinks every day while using this product
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take more or for a longer time than directed

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you have asthma ■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are taking a prescription drug for**

- gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
    - feel faint ■ vomit blood
    - have bloody or black stools
    - have stomach pain that does not get better
  - an allergic reaction occurs. Seek medical help right away.
  - new symptoms occur
  - ringing in the ears or loss of hearing occurs
  - pain gets worse or lasts more than 10 days
  - redness or swelling is present
- These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

**Other information**

- store at 25°C (77°F), excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

**Drug Facts (continued)**

**Inactive ingredients** cherry flavor, corn starch, D&C red #27 aluminum lake, D&C red #30 aluminum lake, dextrates, sodium saccharin

**Questions or comments?** 1-800-426-9391

\*Walgreens Pharmacist Survey Study, November 2010.

†This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Bayer® Chewable Low Dose Aspirin.  
50844 REV0613A39707

ORG112-F\_R01

*Walgreens*  
*Way to Well*  
Convenient

Support local events wellness with Walgreens.  
With each purchase of Walgreens Brand Health & Wellness Products, 1¢ goes to the Walgreens Way to Well fund, up to \$1 million annually. Visit walgreens.com/waytowell

03  
DSC

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

ITEM 535011  
DISTRIBUTED BY: WALGREEN CO.  
200 WILMOT RD, DEERFIELD, IL 60015  
100% SATISFACTION GUARANTEED  
walgreens.com  
©2012 Walgreen Co.



Walgreens 44-397

**LOW DOSE ASPIRIN**

aspirin tablet, chewable

**Product Information**

**Product Type**

HUMAN OTC DRUG

**Item Code (Source)**

NDC:0363-0397

**Route of Administration**

ORAL

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

**Inactive Ingredients**

Ingredient Name	Strength
D&C RED NO. 27 (UNII: 2LRS185U6K)	
D&C RED NO. 30 (UNII: 2S42T2808B)	
DEXTRATES (UNII: G263M44RU)	

**Product Characteristics**

Color	PINK	Score	no score
Shape	ROUND	Size	8mm
Flavor	CHERRY	Imprint Code	44;397
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0397-07	3 in 1 CARTON	09/04/2001	
1		36 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part343	09/04/2001	

**Labeler** - Walgreen Company (008965063)**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(0363-0397)

**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	MANUFACTURE(0363-0397)