

LIL DRUG STORE ANTI-DIARRHEAL- loperamide hydrochloride tablet
Lil' Drug Store Products, Inc.

Anti-Diarrheal, Lil' Drug Store ®

Drug Facts

Active ingredient (in each caplet)

Loperamide HCl, USP 2 mg

Purpose

Anti-diarrheal

Use

Controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert

Do not use if you have ever had a rash or other allergic reaction to Loperamide HCl.

Heart alert

Taking more than directed can cause serious heart problems or death.

Do not use if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise use age

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9 - 11 years (60 - 95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6 - 8 years (48 - 59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2 - 5 years (34 - 47 lbs)	ask a doctor
children under 2 (up to 33 lbs)	do not use

Other information

- store between 68-77°F (20-25°C)

Inactive ingredients

anhydrous lactose, croscarmellose sodium, crospovidone, D&C Yellow no. 10 aluminum lake, FD&C Blue no. 1 aluminum lake, hydrogenated vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch

Questions or comments?

call toll-free **1-877-507-6516 (M-F 8AM - 4:30PM CST)**

PRINCIPAL DISPLAY PANEL - 2 mg Caplet Pouch Carton

QUALITY

★ GUARANTEED ★

Compare to the Active
Ingredient in Imodium® A-D*

Anti-
Diarrheal

on
the
Go
PACKS

See New Warning & Directions

Anti-Diarrheal

Loperamide HCl Tablets, USP 2 mg

- Controls the Symptoms of Diarrhea

4
Caplets**
2 POUCHES, 2 CAPLETS PER POUCH

Lil'
Drug Store®

Compare to the Active Ingredient in Imodium® A-D

Anti-Diarrheal

on the **Go** PACKS

See New Warning & Directions

Anti-Diarrheal

Loperamide HCl Tablets, USP 2 mg

- Controls the Symptoms of Diarrhea

4 Caplets**

2 POUCHES, 2 CAPLETS PER POUCH

Lil' Drug Store

4 Caplets**

Anti-Diarrheal

DO NOT USE IF INNER POUCH OR CARTON IS OPEN OR DAMAGED!

Drug Facts

Active ingredient (in each caplet)	Purpose
Loperamide HCl, USP 2 mg	Anti-diarrheal

Use Controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to Loperamide HCl.

Heart alert: Taking more than directed can cause serious heart problems or death.

Do not use if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

LIFT PANEL FOR CONTINUED DRUG FACTS

Drug Facts (continued)

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- find right dose on chart. If possible, use weight to dose; otherwise use age

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9 - 11 years (60 - 95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6 - 8 years (48 - 59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2 - 5 years (34 - 47 lbs)	ask a doctor
children under 2 (up to 33 lbs)	do not use

4 Caplets**

Anti-Diarrheal

Anti-Diarrheal

Drug Facts (continued)

Other information

- store between 68-77°F (20-25°C)

Inactive ingredients

anhydrous lactose, croscarmellose sodium, crospovidone, D&C Yellow no. 10 aluminum lake, FD&C Blue no. 1 aluminum lake, hydrogenated vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch

Questions or comments?

call toll-free 1-877-507-6516 (M-F 8AM - 4:30PM CST)

Product manufactured for: Lil' Drug Store Products, Inc.
9300 Earhart Lane SW, Cedar Rapids, IA 52404
www.lildrugstore.com 97592C-US-05-19

*This product is not manufactured or distributed by McNeil Consumer Healthcare. Lil' Drug Store Products, Inc. does not own the Imodium® trademark.

Each caplet (**capsule-shaped tablet) contains Loperamide HCl, USP 2 mg

LOT

EXP

8 26529291516671993

4 Caplets**

Anti-Diarrheal Dispens-It, 25pc, Lil' Drug Store [®]

COMPARE TO THE ACTIVE INGREDIENT IN

Imodium [®] A-D*

Anti-Diarrheal

Loperamide HCl Tablets, USP 2 mg

- Controls the Symptoms

of Diarrhea

[tablet image]

Actual Size

This product not for households with small children.

1 CAPLET PER POUCH

[Lil' Drug Store [®] logo]

Drug Facts

Active ingredient (in each caplet)	Purpose
Loperamide HCl, USP 2 mg	Anti-diarrheal

Use
Controls symptoms of diarrhea, including travelers' diarrhea.

Warnings
Allergy alert: Do not use if you have ever had a rash or other allergic reaction to Loperamide HCl. Heart alert: Taking more than directed can cause serious heart problems or death. Do not use if you have bloody or black stool.

Ask a doctor before use if you have:
 • a history of liver disease
 • a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

Stop use and ask a doctor if:
 • symptoms get worse
 • diarrhea lasts for more than 2 days
 • you get abdominal swelling or bloating. These may be signs of a serious condition.

Directions
 • drink plenty of clear fluids to help prevent dehydration caused by diarrhea
 • first: 1 caplet after the first loose stool; if possible, use weight to dose; otherwise use age

Age Group	Dosage
adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool, but no more than 4 caplets in 24 hours
children 9-11 years (80-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool, but no more than 3 caplets in 24 hours
children 6-8 years (40-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool, but no more than 2 caplets in 24 hours
children 2-5 years (24-47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information • store between 68-77° (20-25°C)

Inactive ingredients • anhydrous lactose, croscarmellose sodium, croscopolone, DSG, hollow no. 10 aluminum foil, FD-30, SDC no. 1, aluminum foil, hydroxyethyl cellulose, polyethylene glycol, powdered cellulose, pregelatinized starch

Questions or comments?
call 1-877-507-6516 9A-F (M-F 9AM - 6:30PM CST)

* This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division, distributor of Imodium® A-D.

Packaged and distributed by: Lil' Drug Store Products, Inc. 5900 Earthlink Lane SW, Cedar Rapids, IA 52404

Date Stamp Area
No Varnish

64590-US-55-23

1 CAPLET PER POUCH

Lil' Drug Store

1 CAPLET PER POUCH

Anti-Diarrheal Dispens-It, 30pc, Lil' Drug Store [®]

COMPARE TO THE ACTIVE INGREDIENT IN

Imodium [®] A-D*

Anti-Diarrheal

Loperamide HCl Tablets, USP 2 mg

- Controls the Symptoms

of Diarrhea

[tablet image]

Actual Size

This product not for households with small children.

1 CAPLET PER POUCH

[Lil' Drug Store® logo]



LIL DRUG STORE ANTI-DIARRHEAL

loperamide hydrochloride tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66715-9859
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg
---	--------------------------	------

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE (15 MPA.S AT 5%) (UNII: 68401960MK)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
STARCH, CORN (UNII: O8232NY3SJ)	
GLYCERYL TRISTEARATE (UNII: P6OCJ2551R)	

Product Characteristics

Color	green	Score	2 pieces
Shape	OVAL	Size	10mm
Flavor		Imprint Code	123
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-9859-2	1 in 1 CARTON	09/01/2014	
1		2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:66715-9859-7	50 in 1 CARTON	06/20/2016	09/12/2021
2		2 in 1 POUCH; Type 0: Not a Combination Product		
3	NDC:66715-9859-5	25 in 1 BOX, UNIT-DOSE	05/06/2022	
3		2 in 1 POUCH; Type 0: Not a Combination Product		
4	NDC:66715-9859-6	30 in 1 BOX, UNIT-DOSE	05/06/2022	
4		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074091	09/14/2012	

Revised: 11/2024

Lil' Drug Store Products, Inc.