

IRON HYDROGENATED DEXTRAN- iron injection
MWI Veterinary Supply, Inc.

IRON HYDROGENATED DEXTRAN INJECTION

HEMATINIC

For Prevention and Treatment of Iron Deficiency Anemia in Baby Pigs

NOT FOR HUMAN USE

KEEP OUT OF REACH OF CHILDREN

Approved by FDA under NADA # 138-255

DESCRIPTION

IRON HYDROGENATED DEXTRAN INJECTION is a sterile solution containing 100 mg of elemental iron per mL, stabilized with a low molecular weight hydrogenated dextran and 0.5% phenol as a preservative.

INDICATIONS

For the prevention and treatment of anemia due to iron deficiency in baby pigs.

Iron deficiency anemia occurs commonly in the suckling pig, often within the first few days following birth. As body size and blood volume increase rapidly from the first few days following birth, hemoglobin levels in the blood fall due to diminishing iron reserves which cannot be replaced adequately from iron in the sow's milk. Thus, the need for a readily available outside source of iron is apparent. Without this additional iron many newborn pigs may develop anemia within the first few days of life. Anemia retards growth and lowers the body defenses against disease; scours, pneumonia, and other conditions may develop. Death from anemia or infections may occur, sometimes destroying entire litters.

DOSAGE

For intramuscular injection only.


For **Prevention** of iron deficiency anemia, administer 1 mL (100 mg iron) at 2 to 4 days of age.

For **Treatment** of iron deficiency anemia, administer 1 mL (100 mg iron). Treatment may be repeated in 10 days.

DIRECTIONS FOR USE

Disinfect rubber stopper of vial and injection site with a cotton ball moistened with rubbing alcohol or other suitable disinfectant. Use a small needle (20 gauge, 5/8 inch) and syringe that have been sterilized by boiling in water for 20 minutes. Injection should be made intramuscularly into the back of the ham (see diagram below). Apply tension to the skin at the injection site by pulling skin and fat downward with pressure from the thumb or fingers. Inject into the muscle to a depth of approximately 1/2 inch. Use a

clean, sterile needle and syringe for each pig to avoid spreading infection.



PRECAUTIONS

If disease or abnormal condition is suspected, consult veterinarian before administration.

SIDE EFFECTS

Occasionally pigs may exhibit a reaction to iron injection, clinically characterized by prostration and muscular weakness. In extreme cases, death may result.

NOTICE: The intramuscular injection of organic iron preparations into pigs beyond 4 weeks of age may cause staining of muscle tissue.

HOW SUPPLIED

100 mL multiple dose vials.

Store at controlled room temperature between 15° and 30°C (59° - 86°F).

Protect from freezing.

TAKE TIME OBSERVE LABEL DIRECTIONS

See accompanying leaflet for additional information.

INDICATIONS: For the prevention and treatment of anemia due to iron deficiency in baby pigs.

DOSAGE AND ADMINISTRATION: For intramuscular injection only.


Prevention: 1 mL (100 mg iron) at 2 to 4 days of age.

Treatment: 1 mL (100 mg iron). Treatment may be repeated in 10 days.

NOTICE: Organic iron preparations injected intramuscularly into pigs beyond 4 weeks of age may cause staining of muscle tissue.


See insert for additional information.

Distributed by: MWI
Boise, ID 83705 www.VetOne.net

TAKE TIME  **OBSERVE LABEL DIRECTIONS**

LOT NO.: EXP. DATE:

100 mL

OPEN HERE 

VETone
Iron Hydrogenated Dextran
Injection

HEMATINIC
For Prevention and Treatment of Iron Deficiency Anemia in Baby Pigs

**NOT FOR HUMAN USE
KEEP OUT OF REACH OF CHILDREN**


Approved by FDA under NADA # 138-255

V1 510211
Net Contents: 100 mL (3.4 fl oz)

Each mL of sterile solution contains 100 mg of elemental iron stabilized with a low molecular weight hydrogenated dextran and 0.5% phenol as a preservative.

Store between 15° and 30°C (59° - 86°F).
Protect from freezing.
Use within 18 months of first puncture and puncture a maximum of 200 times.

Contact Sparhawk Laboratories, Inc. at 1-800-888-6368. To report side effects, contact Sparhawk Laboratories, Inc. at 1-800-888-6368. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS (1-888-332-8387) or at <http://www.fda.gov/reportanimalae>.
1-0900-04 Rev. 01/23



3 13985 00893 1

PRECAUTIONS: If disease or abnormal condition is suspected, consult veterinarian before administration.

SIDE EFFECTS: Occasionally pigs may exhibit a reaction to iron injection, clinically characterized by prostration and muscular weakness. In extreme cases, death may result.

NOTICE: The intramuscular injection of organic iron preparations into pigs beyond 4 weeks of age may cause staining of muscle tissue.

HOW SUPPLIED: 100 mL and 250 mL multiple dose vials.

TAKE TIME  **OBSERVE LABEL DIRECTIONS**


Distributed by: MWI
Boise, ID 83705 www.VetOne.net
1-0900-04 Iss. 02/22

5

Each mL of sterile solution contains 100 mg of elemental iron stabilized with a low molecular weight hydrogenated dextran and 0.5% phenol as a preservative.

Store between 15° and 30°C (59° - 86°F).
Protect from freezing.
Use within 18 months of first puncture and puncture a maximum of 200 times.

Contact Sparhawk Laboratories, Inc. at 1-800-888-6368. To report side effects, contact Sparhawk Laboratories, Inc. at 1-800-888-6368. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS (1-888-332-8387) or at <http://www.fda.gov/reportanimalae>.


3 13985 00893 1

VETone
Iron Hydrogenated Dextran
Injection

HEMATINIC
For Prevention and Treatment of Iron Deficiency Anemia in Baby Pigs

DESCRIPTION: Iron Hydrogenated Dextran Injection is a sterile solution containing 100 mg of elemental iron per mL, stabilized with a low molecular weight hydrogenated dextran and 0.5% phenol as a preservative.

INDICATIONS: For the prevention and treatment of anemia due to iron deficiency in baby pigs.

1

Iron deficiency anemia occurs commonly in the suckling pig, often within the first few days following birth. As body size and blood volume increase rapidly from the first few days following birth, hemoglobin levels in the blood fall due to diminishing iron reserves which cannot be replaced adequately from iron in the sow's milk. Thus, the need for a readily available outside source of iron is apparent. Without this additional iron many newborn pigs may develop anemia within the first few days of life. Anemia retards growth and lowers the body defenses against disease, scours, pneumonia, and other conditions may develop. Death from anemia or infections may occur, sometimes destroying entire litters.

2

DOSAGE: For intramuscular injection only.


For **Prevention** of iron deficiency anemia, administer 1 mL (100 mg iron) at 2 to 4 days of age.

For **Treatment** of iron deficiency anemia, administer 1 mL (100 mg iron). Treatment may be repeated in 10 days.

DIRECTIONS FOR USE: Disinfect rubber stopper of vial and injection site with a cotton ball moistened with rubbing alcohol or other suitable disinfectant. Use a small needle (20 gauge, 5/8 inch) and syringe that have been sterilized by boiling in water for 20 minutes.

3

Injection should be made intramuscularly into the back of the ham (see diagram below). Apply tension to the skin at the injection site by pulling skin and fat downward with pressure from the thumb or fingers. Inject into the muscle to a depth of approximately 1/2 inch. Use a clean, sterile needle and syringe for each pig to avoid spreading infection.



4

IRON HYDROGENATED DEXTRAN

iron injection

Product Information			
Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:13985-520
Route of Administration	INTRAMUSCULAR		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IRON (UNII: E1UOL152H7) (IRON - UNII:E1UOL152H7)	IRON	100 mg in 1 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:13985-520-02	100 mL in 1 VIAL		
2	NDC:13985-520-05	250 mL in 1 VIAL		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA138255	11/19/2011	

Labeler - MWI Veterinary Supply, Inc. (019926120)

Revised: 10/2023

MWI Veterinary Supply, Inc.