

TRIAMCINOLONE ACETONIDE - triamcinolone acetonide paste
AvKARE



TRIAMCINOLONE ACETONIDE DENTAL PASTE USP, 0.1%

Rx only

Rx only

DESCRIPTION

Triamcinolone Acetonide Dental Paste USP, 0.1% contains the corticosteroid triamcinolone acetonide in an adhesive vehicle suitable for application to oral tissues. Triamcinolone acetonide is designated chemically as 9-fluoro-11 β , 16 α , 17, 21-tetrahydroxypregna-1, 4-diene-3, 20-dione cyclic 16,17-acetal with acetone. The structural formula of triamcinolone acetonide is as follows:



Each gram of Triamcinolone Acetonide Dental Paste USP, 0.1% contains 1 mg triamcinolone acetonide in a dental paste containing gelatin, pectin, cream flavor, vanilla flavor and carboxymethylcellulose sodium in Plasticized Hydrocarbon Gel, a polyethylene and mineral oil gel base.

CLINICAL PHARMACOLOGY

Like other topical corticosteroids, triamcinolone acetonide has anti-inflammatory, antipruritic, and vasoconstrictive properties. The mechanism of the anti-inflammatory activity of the topical steroids, in general, is unclear. However, corticosteroids are thought to act by the induction of phospholipase A₂ inhibitory proteins, collectively called lipocortins. It is postulated that these proteins control the biosynthesis of potent mediators of inflammation such as prostaglandins and leukotrienes by inhibiting the release of their common precursor, arachidonic acid. Arachidonic acid is released from membrane phospholipids by phospholipase A₂.

Pharmacokinetics

The extent of absorption through the oral mucosa is determined by multiple factors including the

vehicle, the integrity of the mucosal barrier, the duration of therapy, and the presence of inflammation and/or other disease processes. Once absorbed through the mucous membranes, the disposition of corticosteroids is similar to that of systemically administered corticosteroids. Corticosteroids are bound to the plasma proteins in varying degrees. Corticosteroids are metabolized primarily in the liver and are then excreted by the kidneys; some corticosteroids and their metabolites are also excreted into the bile.

INDICATIONS AND USAGE

Triamcinolone Acetonide Dental Paste USP, 0.1% is indicated for adjunctive treatment and for the temporary relief of symptoms associated with oral inflammatory lesions and ulcerative lesions resulting from trauma.

CONTRAINDICATIONS

Triamcinolone Acetonide Dental Paste USP, 0.1% is contraindicated in those patients with a history of hypersensitivity to any of the components of the preparations; it is also contraindicated in the presence of fungal, viral, or bacterial infections of the mouth or throat.

PRECAUTIONS

General:

Triamcinolone Acetonide Dental Paste USP, 0.1% may cause local adverse reactions. If irritation develops, Triamcinolone Acetonide Dental Paste USP, 0.1% should be discontinued and appropriate therapy instituted. Allergic contact sensitization with corticosteroids is usually diagnosed by observing failure to heal rather than noting a clinical exacerbation as with most topical products not containing corticosteroids. Such an observation should be corroborated with appropriate diagnostic patch testing.

If concomitant mucosal infections are present or develop, an appropriate antifungal or antibacterial agent should be used. If a favorable response does not occur promptly, use of Triamcinolone Acetonide Dental Paste USP, 0.1% should be discontinued until the infection has been adequately controlled.

If significant regeneration or repair of oral tissues has not occurred in seven days, additional investigation into the etiology of the oral lesion is advised.

Systemic absorption of topical corticosteroids has produced reversible hypothalamic-pituitary-adrenal (HPA) axis suppression, manifestations of Cushing's syndrome, hyperglycemia, glucosuria, and other adverse effects known to occur with parenterally-administered steroid preparations; therefore, it may be advisable to periodically evaluate patients on prolonged therapy with corticosteroid-containing dental pastes for evidence of HPA axis suppression (see **PRECAUTIONS, Laboratory Tests**). If HPA axis suppression is noted, an attempt should be made to withdraw the drug or to reduce the frequency of application. Recovery of HPA axis function is generally prompt and complete upon discontinuation of therapy.

Information for the Patient: Patients using topical corticosteroids should receive the following information and instructions:

This medication is to be used as directed by the physician or dentist. It is for oral use only; it is not intended for ophthalmic or dermatological use.

Patients should be advised not to use this medication for any disorder other than for which it was prescribed.

Patients should report any signs of adverse reactions.

As with other corticosteroids, therapy should be discontinued when control is achieved. If no improvement is seen within 2 weeks, contact the physician or dentist.

Laboratory Tests

A urinary free cortisol test and ACTH stimulation test may be helpful in evaluating HPA axis suppression.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Animal studies have not been performed to evaluate triamcinolone acetonide for potential to induce carcinogenesis, mutagenesis, or impairment of fertility.

Pregnancy Category C: Teratogenic effects. Triamcinolone acetonide has been shown to induce teratogenic effects in several species. In mice and rabbits, triamcinolone acetonide induced an increased incidence of cleft palate at dosages of approximately 120 mcg/kg/day and 24 mcg/kg/day, respectively (approximately 12 times and 10 times the amount in a typical daily human dose of Triamcinolone Acetonide Dental Paste USP, 0.1% when compared following normalization of the data on the basis of body surface area estimates, respectively). In monkeys, triamcinolone acetonide induced cranial skeletal malformations at the lowest dosage studied (500 mcg/kg/day), which was approximately 200 times the amount in a typical daily human dose of Triamcinolone Acetonide Dental Paste USP, 0.1% when compared following normalization of the data on the basis of body surface area estimates. There are no adequate and well-controlled studies in pregnant women. However, a retrospective analysis of birth defects among children born to mothers that used drugs of the same class as Triamcinolone Acetonide Dental Paste USP, 0.1% (corticosteroids) during pregnancy found an approximately 3 times increased incidence of cleft palate. Triamcinolone Acetonide Dental Paste USP, 0.1% should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nursing Mothers

It is not known whether oral application of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in breast milk. Caution should be exercised when corticosteroid-containing dental pastes are prescribed for a nursing woman.

Pediatric Use

The safety and efficacy of Triamcinolone Acetonide Dental Paste USP, 0.1% in children is unknown. Pediatric patients may demonstrate greater susceptibility to topical corticosteroid-induced HPA axis suppression and Cushing's Syndrome than mature patients because of a larger skin surface area to body weight ratio. Administration of corticosteroid-containing dental pastes to children should be limited to the least amount compatible with an effective therapeutic regimen. Chronic corticosteroid therapy may interfere with the growth and development of children.

Geriatric Use

Clinical studies of Triamcinolone Acetonide Dental Paste USP, 0.1% did not include sufficient numbers of subjects age 65 and older to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients.

ADVERSE REACTIONS

The following local adverse reactions may occur with corticosteroid-containing dental pastes: burning, itching, irritation, dryness, blistering or peeling not present prior to therapy, perioral dermatitis, allergic contact dermatitis, maceration of the oral mucosa, secondary infection, and atrophy of the oral mucosa.

Also, see **PRECAUTIONS** for potential effects of systemic absorption.

DOSAGE AND ADMINISTRATION

Press a small dab (about 1/4 inch) to the lesion until a thin film develops. A larger quantity may be required for coverage of some lesions. For optimal results use only enough to coat the lesion with a

thin film. Do not rub in. Attempting to spread this preparation may result in granular, gritty sensation and cause it to crumble. After application, however, a smooth, slippery film develops.

The preparation should be applied at bedtime to permit steroid contact with the lesion throughout the night. Depending on the severity of symptoms, it may be necessary to apply the preparation two or three times a day, preferably after meals. If significant repair or regeneration has not occurred in seven days, further investigation is advisable.

HOW SUPPLIED

Triamcinolone Acetonide Dental Paste USP, 0.1%
Tubes containing 5 g of dental paste
NDC 42291-840-05

Storage

Keep tightly closed. Store at 20° - 25° C (68° - 77° F) excursion permitted to 15° - 30° C (59° - 86° F) [See USP Controlled Room Temperature].

Distributed by:

AvKARE, Inc.
Pulaski, TN 38478

Manufactured by:

Lyne Laboratories, Inc.
Brockton, MA 02301

RO-04/12

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

AvKARE **NDC 42291-840-05**

Triamcinolone Acetonide
Dental Paste USP, 0.1%

5g Rx Only
 Read all sides

5g
Triamcinolone
Acetonide
Dental Paste
USP, 0.1%

EACH GRAM CONTAINS: 1mg triamcinolone acetonide (0.1%) in an emollient dental paste composed of gelatin, pectin and carboxymethylcellulose sodium in a plasticized hydrocarbon gel (a polyethylene and mineral oil gel base).

FOR TOPICAL USE IN ADJUNCTIVE TREATMENT OF ORAL LESIONS.

USUAL DOSAGE: Apply a thin film to affected area, preferably at bedtime, See Insert.
KEEP TIGHTLY CLOSED.

Store at 20o-25oC (68o-77oF) excursions permitted to 15o-30oC (59o-86oF) [See USP Controlled Room Temperature]. For lot number and expiry date see flap of carton or crimp of tube.

AvKARE NDC 42291-840-05

Triamcinolone Acetonide
Dental Paste USP, 0.1%

5g Rx Only
Read all sides

Manufactured by:
Lyne Laboratories, Inc.,
Brockton, MA 02301

Distributed by:
AvKARE, Inc.,
Pulaski, TN 38478

3 42291 84005 8

R1 - 10/16





NDC 42291-840-05

Triamcinolone Acetonide Dental Paste USP, 0.1%

Rx only

Net Wt. 5g

EACH GRAM CONTAINS: 1 mg triamcinolone acetonide (0.1%) in an emollient dental paste composed of gelatin, pectin and carboxymethylcellulose sodium in plasticized hydrocarbon gel (a polyethylene and mineral oil gel base).

FOR TOPICAL USE IN ADJUNCTIVE TREATMENT OF ORAL LESIONS.

USUAL DOSAGE: Apply a thin film to affected area, preferably at bedtime. See Insert.

READ ALL SIDES. KEEP TIGHTLY CLOSED.

Store at 20°-25°C (68°-77°F) excursion permitted to 15°-30°C (59°-86°F) [See USP Controlled Room Temperature].

For lot number and expiry date see crimp.

Mfd. by: Lyne Laboratories, Inc., Brockton, MA 02301

Distributed by: AvKARE, Inc., Pulaski, TN 38478

R0 - 04/12



TRIAMCINOLONE ACETONIDE

triamcinolone acetonide paste

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:42291-840
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TRIAMCINOLONE ACETONIDE (UNII: F446C597KA) (TRIAMCINOLONE ACETONIDE - UNII:F446C597KA)	TRIAMCINOLONE ACETONIDE	1 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
GELATIN (UNII: 2G86QN327L)	
PECTIN, CITRUS (UNII: 47EQ08LE7H)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)	

Product Characteristics

Color		Score	
--------------	--	--------------	--

Shape		Size	
Flavor	VANILLA (Vanilla)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42291-840-05	48 in 1 BOX	06/07/2012	
1		1 in 1 CARTON		
1		5 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040771	06/07/2012	

Labeler - AvKARE (796560394)

Revised: 1/2020

AvKARE