

VICKS NYQUIL COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, and doxylamine succinate liquid
The Procter & Gamble Manufacturing Company

VICKS[®] NyQuil[™] COLD & FLU

Drug Facts

Active ingredients (in each 30 mL dose cup)

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, see a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 30 mL dose cup contains:**sodium 23 mg
- store at no greater than 25°C and do not refrigerate

Inactive ingredients

alcohol, citric acid, D&C Yellow No. 10, FD&C Green No. 3, FD&C Yellow No. 6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions?

1-800-362-1683

TAMPER EVIDENT: Do not use if printed shrinkband is broken or missing.

DIST. BY

PROCTER & GAMBLE,

CINCINNATI, OH 45202

PRINCIPAL DISPLAY PANEL - 236 ml Bottle Label

VICKS ®

NyQuil ™

COLD & FLU

Nighttime Relief



Acetaminophen, Doxylamine Succinate, Dextromethorphan HBr

Headache, Fever, Sore Throat, Minor Aches & Pains
Sneezing, Runny Nose
Cough

Nighttime Relief

Alcohol 10%

8 FL OZ (236 ml)



VICKS NYQUIL COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, and doxylamine succinate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69423-789
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	

FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
XANTHAN GUM (UNII: TTV12P4NEE)	
GLYCERIN (UNII: PDC6A3C0OX)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	green	Score	
Shape		Size	
Flavor	ANISE, MENTHOL	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69423-789-08	236 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/01/2022	
2	NDC:69423-789-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/01/2022	
3	NDC:69423-789-24	2 in 1 PACKAGE, COMBINATION	01/01/2022	
3	NDC:69423-789-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/01/2022	

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Revised: 10/2023

The Procter & Gamble Manufacturing Company