

**MUCINEX FAST MAX LIQUID COLD AND FLU KICKSTART AND MUCINEX NIGHTSHIFT KICKSTART- acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hci, triprolidine hci
RB Health (US) LLC**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Mucinex Fast-Max Kickstart Severe Cold & Flu and Night Time Cold & Flu

Drug Facts

KICKSTART SEVERE COLD & FLU:

Acetaminophen 650mg.....Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....Cough suppressant
Guaifenesin 400 mg.....Expectorant
Phenylephrine HCl 10mg.....Nasal decongestant

NIGHT TIME SEVERE COLD & FLU:

Acetaminophen 650mg.....Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....Cough suppressant
Phenylephrine HCl 10mg.....Nasal decongestant
Triprolidine HCl 2.5 mg.....Antihistamine

Uses

KICKSTART SEVERE COLD & FLU:

temporarily relieves these common cold and flu symptoms:

- cough
- nasal congestion
- minor aches and pains
- sore throat
- headache
- stuffy nose
- sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

NIGHT TIME SEVERE COLD & FLU:

temporarily relieves these common cold and flu symptoms:

- cough
- nasal congestion

- minor aches and pains
- sore throat
- headache
- sinus congestion and pressure
- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

KICKSTART SEVERE COLD & FLU:

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease

- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

NIGHT TIME SEVERE COLD & FLU:

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription).
- If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts

These could be signs of a serious condition.

Directions

KICKSTART SEVERE COLD & FLU

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

NIGHT TIME SEVERE COLD & FLU

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other Information

KICKSTART SEVERE COLD & FLU

- each 20 mL contains: sodium 12 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

NIGHT TIME SEVERE COLD & FLU

- each 20 mL contains: sodium 16 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

KICKSTART SEVERE COLD & FLU

ammonium glycyrrhizate, anhydrous citric acid, D&C yellow no. 10, edetate disodium, FD&C blue no. 1, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

NIGHT TIME SEVERE COLD & FLU

ammonium glycyrrhizate, anhydrous citric acid, ascorbic acid, edetate disodium, FD&C blue no. 1,

FD&C red no. 40, flavors, glycerin (soy), propylene glycol, sodium benzoate, sorbitol, sucralose, triacetin, triethyl citrate, water, xanthan gum

1-866-MUCINEX
(1-866-682-4639)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Purposes

KICKSTART SEVERE COLD & FLU:

Dextromethorphan HBr 20 mg.....Cough suppressant
Guaifenesin 400 mg.....Expectorant
Phenylephrine HCl 10mg.....Nasal decongestant

NIGHT TIME SEVERE COLD & FLU:

Dextromethorphan HBr 20 mg.....Cough suppressant
Phenylephrine HCl 10mg.....Nasal decongestant
Triprolidine HCl 2.5 mg.....Antihistamine

NDC 72854-139

MUCINEX FAST-MAX Kickstart + Nighttime Cold & Flu Pack

Kickstart Severe Cold & Flu:

Acetaminophen – Pain Reliever/Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Guaifenesin – Expectorant • Phenylephrine HCl – Nasal Decongestant

Nighttime Severe Cold & Flu:

Acetaminophen – Pain Reliever/Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Phenylephrine HCl – Nasal Decongestant • Triprolidine HCl – Antihistamine

FOR

AGES 12+

TWO – 6 FL OZ (180 mL) bottles

TOTAL – 12 FL OZ (360 mL)

A KICKSTART FOR YOUR DAY

MAXIMUM STRENGTH

POWERFUL RELIEF FOR YOUR NIGHT



Mucinex
FAST-MAX
KICKSTART

Mucinex
FAST-MAX
NIGHT-TIME



POWERFUL MULTI-SYMPTOM COLD & FLU RELIEF

ALL IN ONE SEVERE COLD & FLU RELIEF » DAY OR NIGHT

KICKSTART SEVERE COLD & FLU (cont'd)

Drug Facts (continued)

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients

ammonium glycyrrhizate, anhydrous citric acid, D&C yellow no. 10, edetate disodium, FD&C blue no. 1, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions? 1-866-MUCINEX (1-866-682-4639)

*FAST-MAX KICKSTART SEVERE COLD & FLU helps to relieve these symptoms day or night; FAST-MAX NIGHT TIME SEVERE COLD & FLU helps to relieve these symptoms at night

Do not take Mucinex Fast-Max Kickstart Severe Cold & Flu and Mucinex Fast-Max Night Time Severe Cold & Flu at the same time. Always wait at least 4 hours before taking another dose of Mucinex Liquid.

TAKE ONLY AS DIRECTED

Maximum Strength per 4-hour dose



FOR AGES 12+

MENTHOL FLAVORED LIQUID

TWO - 6 FL OZ (180 mL) bottles

RELIEF FOR YOUR NIGHT

TOTAL - 12 FL OZ (360 mL)

NIGHT-TIME SEVERE COLD & FLU (cont'd)

Drug Facts (continued)

- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 16 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients

ammonium glycyrrhizate, anhydrous citric acid, ascorbic acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin (scl), propylene glycol, sodium benzoate, sorbitol, sucralose, tetracain, triethyl citrate, water, xanthan gum

Questions? 1-866-MUCINEX (1-866-682-4639)

POWERFUL RELIEF FOR YOUR NIGHT
+
A KICKSTART FOR YOUR DAY



Dist. by: RB Health (US)
Parsippany, NJ 07054-8224
© 2023 RB Health
3259878 040523

NIGHT TIME SEVERE COLD & FLU

Drug Facts

Active Ingredients (In each 20 mL)

Purposes

| | |
|----------------------------|-----------------------------|
| Acetaminophen 650 mg | Pain reliever/fever reducer |
| Dextromethorphan HBr 20 mg | Cough suppressant |
| Phenylephrine HCl 10 mg | Nasal decongestant |
| Triprolidine HCl 2.5 mg | Antihistamine |

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - sinus congestion and pressure
 - runny nose
 - sneezing
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)
- liver disease
- heart disease
- thyroid disease
- glaucoma

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product do not use more than directed

- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days

Tamper evident: Do not use if neckband on bottle cap is broken or missing.

KICKSTART SEVERE COLD & FLU

Drug Facts

Active Ingredients (In each 20 mL)

Purposes

| | |
|----------------------------|-----------------------------|
| Acetaminophen 650 mg | Pain reliever/fever reducer |
| Dextromethorphan HBr 20 mg | Cough suppressant |
| Guaifenesin 400 mg | Expectorant |
| Phenylephrine HCl 10 mg | Nasal decongestant |

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - stuffy nose
 - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)
- liver disease
- heart disease
- thyroid disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

MAXIMUM STRENGTH

NDC 72854-138-66

3259831

030823

Mucinex[®]

FAST[®]MAX[®]

KICKSTART SEVERE COLD & FLU

Acetaminophen » Pain Reliever/Fever Reducer

Dextromethorphan HBr » Cough Suppressant

Guaifenesin » Expectorant

Phenylephrine HCl » Nasal Decongestant

**COMEBACK
WITH A
KICK**



MENTHOL FLAVORED LIQUID

6 FL OZ (180 mL)

FOR AGES 12+

Mucinex Mucinex Mucinex Mucinex Mucinex
Mucinex Mucinex Mucinex Mucinex Mucinex
Mucinex Mucinex Mucinex Mucinex Mucinex
Mucinex Mucinex Mucinex Mucinex Mucinex

Mucinex Mucinex Mucinex Mucinex Mucinex
Mucinex Mucinex Mucinex Mucinex Mucinex
Mucinex Mucinex Mucinex Mucinex Mucinex
Mucinex Mucinex Mucinex Mucinex Mucinex



MUCINEX FAST MAX LIQUID COLD AND FLU KICKSTART AND MUCINEX NIGHTSHIFT KICKSTART

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hci, triprolidine hci kit

Product Information

| | | | |
|---------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:72854-139 |
|---------------------|----------------|---------------------------|---------------|

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------|----------------------|--------------------|
| 1 | NDC:72854-139-26 | 1 in 1 CARTON | 05/01/2023 | |

Quantity of Parts

| Part # | Package Quantity | Total Product Quantity |
|--------|------------------|------------------------|
| Part 1 | 1 BOTTLE | 180 mL |
| Part 2 | 1 BOTTLE | 180 mL |

Part 1 of 2

MUCINEX FAST-MAX KICKSTART SEVERE COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride solution

Product Information

| | |
|-------------------------|---------------|
| Item Code (Source) | NDC:72854-138 |
| Route of Administration | ORAL |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------------------|-----------------|
| PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) | PHENYLEPHRINE HYDROCHLORIDE | 10 mg in 20 mL |
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 20 mg in 20 mL |
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 650 mg in 20 mL |
| GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ) | GUAIFENESIN | 400 mg in 20 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| D&C YELLOW NO. 10 (UNII: 35SW5USQ3G) | |
| FD&C BLUE NO. 1 (UNII: H3R47K3TBD) | |
| SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR) | |
| SORBITOL (UNII: 506T60A25R) | |
| SUCRALOSE (UNII: 96K6UQ3ZD4) | |
| XANTHAN GUM (UNII: TTV12P4NEE) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| SODIUM BENZOATE (UNII: OJ245FE5EU) | |
| WATER (UNII: 059QF0KO0R) | |
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | |
| EDETATE DISODIUM (UNII: 7FLD91C86K) | |
| PROPYL GALLATE (UNII: 8D4SNN7V92) | |

Product Characteristics

| | | | |
|-------|-------|-------|--|
| Color | green | Score | |
| Shape | | Size | |

| | | | |
|-----------------|---------|---------------------|--|
| Flavor | MENTHOL | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:72854-138-66 | 180 mL in 1 BOTTLE; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product) | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part341 | 05/01/2023 | |

Part 2 of 2

MUCINEX NIGHT TIME SEVERE COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride, and triprolidine hydrochloride solution

Product Information

| | |
|--------------------------------|---------------|
| Item Code (Source) | NDC:72854-140 |
| Route of Administration | ORAL |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------------------|-----------------|
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 20 mg in 20 mL |
| PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1W5297W6MV) | PHENYLEPHRINE HYDROCHLORIDE | 10 mg in 20 mL |
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 650 mg in 20 mL |
| TRIPROLIDINE HYDROCHLORIDE (UNII: YAN7R5L890) (TRIPROLIDINE - UNII:2L8T9S52QM) | TRIPROLIDINE HYDROCHLORIDE | 2.5 mg in 20 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| FD&C BLUE NO. 1 (UNII: H3R47K3TBD) | |
| FD&C RED NO. 40 (UNII: WZB9127XOA) | |
| TRIETHYL CITRATE (UNII: 8Z96QXD6UM) | |
| AMMONIUM GLYCYRRHIZATE (UNII: 3VRD35U26C) | |
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | |

| | |
|--|--|
| SODIUM BENZOATE (UNII: OJ245FE5EU) | |
| SORBITOL (UNII: 506T60A25R) | |
| WATER (UNII: 059QF0KO0R) | |
| XANTHAN GUM (UNII: TTV12P4NEE) | |
| ASCORBIC ACID (UNII: PQ6CK8PD0R) | |
| EDETATE DISODIUM (UNII: 7FLD91C86K) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| SUCRALOSE (UNII: 96K6UQ3ZD4) | |
| TRIACETIN (UNII: XHX3C3X673) | |

Product Characteristics

| | | | |
|-----------------|-------|---------------------|--|
| Color | blue | Score | |
| Shape | | Size | |
| Flavor | FRUIT | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:72854-140-66 | 180 mL in 1 BOTTLE; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product) | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part341 | 05/01/2023 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part341 | 05/01/2023 | |

Labeler - RB Health (US) LLC (081049410)