

ANTACID AND ANTIGAS- aluminum hydroxide, magnesium hydroxide, simethicone liquid
Rij Pharmaceutical Corporation

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients (in each 5 mL) (One teaspoonful)

Aluminum Hydroxide (Equi. to Dried Gel, USP) 200 mg

Magnesium Hydroxide 200 mg

Simethicone 20 mg

Purpose

Antacid

Antigas

Uses

relieves

- heartburn
- acid indigestion
- sour stomach
- associated symptoms of gas and upset stomach
- pressure and bloating commonly referred to as gas

Warnings

Do not take more than 16 teaspoonfuls in a 24-hour period or use the maximum dosage for more than 2 weeks. Except under the advise and supervision of a physician.

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if

symptoms last for more than 2 weeks

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center immediately.

Directions

- **shake well before each use**
- Adults/children 12 years and older, take 2 to 4 teaspoons between meals, at bedtime or as directed by a physician
- do not take more than 16 teaspoonsful in 24 hours
- do not use the maximum dosage for more than 2 weeks
- Children under 12 years: consult a physician

Other information

- TAMPER EVIDENT: DO NOT USE IF BREAKAWAY BAND ON CAP IS BROKEN OR MISSING.
- Store at room temperature 15° - 30°C (59° - 86°F)
- keep tightly closed
- protect from freezing
- each teaspoon (5 mL) contains: magnesium 83 mg
- each teaspoon (5 mL) contains: sodium 1 mg
- does not meet USP requirements for preservative effectiveness

Inactive ingredients

flavors, hypromellose, parabens, saccharin, sodium hypochlorite, sorbitol, water and xanthan gum

Principal Display Panel



NDC 53807-538-12
Compare to the Active
Ingredient in Maalox®†

Antacid & Antigas

Magnesia and Alumina
Oral Suspension
with Simethicone

Non-Constipating
Pleasant Tasting



Cherry Flavored

12 FL. OZ (355 mL)

Antacid & Antigas Cherry

Drug Facts

Active Ingredients

(In each 5 ml) (One teaspoonful)

Purpose

Aluminum Hydroxide

(Equi. to Dried Gel, USP) 200 mgAntacid

Magnesium Hydroxide 200 mgAntacid

Simethicone 20 mgAntigas

Uses relieves ■ heartburn ■ acid indigestion ■ sour stomach
■ associated symptoms of gas and upset stomach ■ pressure and
bloating commonly referred to as gas

Warnings

Do not take more than 16 teaspoonfuls in a 24-hour period or use
the maximum dosage for more than 2 weeks. Except under the advise
and supervision of a physician.

Ask a doctor before use if you have

■ kidney disease ■ a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Antacids may interact with certain
prescription drugs.

Stop use and ask a doctor if ■ symptoms last more than 2 weeks

Keep out of reach of children. In case of accidental overdose,
get medical help or contact a Poison Control Center immediately.

Directions ■ shake well before each use

■ Adults/children 12 years and older, take 2 to 4 teaspoons between meals,
at bedtime or as directed by a physician

■ do not take more than 16 teaspoonsful in 24 hours

■ do not use maximum dosage for more than 2 weeks

■ Children under 12 years: consult a physician

Other Information

■ **TAMPER EVIDENT: DO NOT USE IF BREAKAWAY BAND ON CAP IS BROKEN
OR MISSING.**

■ store at room temperature 15°-30°C (59°-86°F)

■ keep tightly closed ■ protect from freezing

■ each teaspoon (5 mL) contains: magnesium 83 mg

■ each teaspoon (5 mL) contains: sodium 1 mg

■ does not meet USP requirements for preservative effectiveness

Inactive ingredients: flavors, hypromellose, parabens, saccharin,
sodium hypochlorite, sorbitol, water and xanthan gum.

Questions? 845-692-5799

†This product is not manufactured or distributed by the owner of the registered trademark Maalox®
All Preferred Plus Pharmacy products are satisfaction guaranteed or your money back
from retailer where purchased.

Manufactured By:
RIJ PHARMACEUTICAL CORP.
40 Commercial Avenue
Middletown, NY 10941
Distributed By:
Kinray Inc.,
Whitestone, NY 11357



LRV.10/12

aluminum hydroxide, magnesium hydroxide, simethicone liquid						
Product Information						
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53807-538			
Route of Administration	ORAL					
Active Ingredient/Active Moiety						
Ingredient Name		Basis of Strength	Strength			
ALUMINUM HYDRO XIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)		ALUMINUM HYDROXIDE	200 mg in 5 mL			
MAGNESIUM HYDRO XIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)		MAGNESIUM HYDROXIDE	200 mg in 5 mL			
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)		DIMETHICONE	20 mg in 5 mL			
Inactive Ingredients						
Ingredient Name			Strength			
HYPROMELLOSES (UNII: 3NXW29V3WO)						
BUTYL PARABEN (UNII: 3QPIIU3FV8)						
SACCHARIN (UNII: FST467XS7D)						
SODIUM HYPOCHLORITE (UNII: DY38VHM5OD)						
SORBITOL (UNII: 506T60A25R)						
PROPYL PARABEN (UNII: Z8IX2SC1OH)						
WATER (UNII: 059QF0KO0R)						
XANTHAN GUM (UNII: TTV12P4NEE)						
Product Characteristics						
Color	WHITE (opaque)	Score				
Shape		Size				
Flavor	CHERRY	Imprint Code				
Contains						
Packaging						
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:53807-538-12	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/16/1999			
Marketing Information						
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date			
OTC monograph final	part332	03/16/1999				

Establishment

Name	Address	ID/FEI	Business Operations
Rij Pharmaceutical Corporation		144679156	manufacture(53807-538)

Revised: 4/2018

Rij Pharmaceutical Corporation