

CHILDRENS PAIN AND FEVER- acetaminophen suspension
CVS PHARMACY, INC

Drug Facts

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily reduces fever
- temporarily relieves minor aches and pains due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this

product

Ask a doctor before use if your child has

liver disease.

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin.

When using this product,

do not exceed recommended dosage (see Overdose warning).

Stop use and ask a doctor if

- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not take more than directed (see Overdose warning)**
- **shake well before using**
- mL= milliliter
- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give more than 5 days unless directed by a doctor
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.

Weight (lb)	Age (year)	Dose (mL)*
under 24	under 2	ask a doctor
24-35	2-3	5 mL
36-47	4-5	7.5 ml
48-59	6-8	10 mL
60-71	9-10	12.5 mL

72-95

11

15 mL

*or as directed by a doctor

Other information

- **each 5 mL contains:** sodium 2 mg
- store between 20-25°C (68-77°F)
- do not refrigerate
- see bottom panel for lot number and expiration date

Inactive ingredients

acesulfame potassium, butylparaben, carboxymethylcellulose sodium, citric acid, FD&C red #40, flavors, glycerin, high fructose corn syrup, microcrystalline cellulose, propylene glycol, purified water, sodium benzoate, sorbitol, xanthan gum

Questions or Comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredient in Children's Tylenol® Oral Suspension**

Oral Suspension

Children's Pain & Fever

ACETAMINOPHEN 160 mg /5 mL

Pain reliever;

Fever reducer

Ages 2 to 11 years

Free of alcohol, aspirin, ibuprofen & gluten

Cherry Flavor

Dosage Cup Included

CHERRY FLAVOR

FL OZ (mL)

IMPORTANT: Please see dosing chart or ask a doctor for your child's correct dose.

**This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Oral Suspension.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

Distributed by: CVS Pharmacy, Inc.
 One CVS Drive, Woonsocket, RI 02895

Product Label

Drug Facts (continued)

- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give more than 5 days unless directed by a doctor
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.

Weight (lb)	Age (year)	Dose (mL)*
under 24	under 2	ask a doctor
24-35	2-3	5 mL
36-47	4-5	7.5 mL
48-59	6-8	10 mL
60-71	9-10	12.5 mL
72-85	11	15 mL

*or as directed by a doctor

Other information

- each 5 mL contains: sodium 2 mg
- store between 20-25°C (68-77°F) ■ do not refrigerate
- see bottom panel for lot number and expiration date

Inactive ingredients: acetaminophen, butylparaben, carboxymethylcellulose sodium, citric acid, FD&C red #40, flavors, glycerin, high fructose corn syrup, microcrystalline cellulose, propylene glycol, purified water, sodium benzoate, sorbitol, xanthan gum

Questions or comments?
 Call 1-877-753-3835 Monday-Friday 9AM-6PM EST

***This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Oral Suspension.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

Distributed by: CVS Pharmacy, Inc.
 One CVS Drive, Woonsocket, RI 02895
 © 2023 CVS/pharmacy
 CVS.com® 1-800-SHOP CVS
 V-19854

100% money back guaranteed.
 CVS.com/returnpolicy

IMPORTANT:
 Please see dosing chart or ask a doctor for your child's correct dose.

Package Contains One Bottle

Actual Size

Compare to the active ingredient in Children's Tylenol® Oral Suspension**

Oral Suspension

NDC 51316-282-04

Children's Pain & Fever

ACETAMINOPHEN
 160 mg / 5 mL

**Pain reliever;
 Fever reducer**

Ages 2 to 11 years

Free of alcohol, aspirin, ibuprofen & gluten

Cherry Flavor
 Dosing Cup Included

4 FL OZ (118 mL)

Actual Bottle Size on Side Panel

Drug Facts

Active Ingredient (in each 5 mL)	Purpose
Acetaminophen 160 mg	Pain reliever/fever reducer

Uses

- temporarily reduces fever
- temporarily relieves minor aches and pains due to
 - the common cold
 - flu
 - headache
- sore throat
- toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dose (see Overdose warning).

Stop use and ask a doctor if

- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not take more than directed (see Overdose warning)
- shake well before using
- mL = milliliter

#207688

0 50428 30779 3

PLD-A231D FC008588

Lot No:
 Exp. Date:

CVS HEALTH Children's Pain and Fever

CHILDRENS PAIN AND FEVER

acetaminophen suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51316-282
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
BUTYLPARABEN (UNII: 3QPI1U3FV8)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KOOR)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51316-282-04	1 in 1 BOX	02/15/2023	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/15/2023	

Labeler - CVS PHARMACY, INC (062312574)