

**EXCHANGE SELECT ACID REDUCER- ranitidine tablet**  
**Army & Air Force Exchange Service**

-----  
**Perrigo Company Acid Reducer Drug Facts**

**Active ingredient (in each tablet)**

Ranitidine 75 mg (as ranitidine hydrochloride 84 mg)

**Purpose**

Acid reducer

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

**Warnings**

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

**Ask a doctor before use if you have**

- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- adults and children 12 years and over:
- to relieve symptoms, swallow 1 tablet with a glass of water
- to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn
- can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

**Other information**

- do not use if printed foil under cap is broken or missing
- store at 20° - 25°C (68° - 77°F)
- avoid excessive heat or humidity
- this product is sugar free

**Inactive ingredients**

colloidal silicon dioxide, croscarmellose sodium, diethyl phthalate, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, titanium dioxide.

**Questions or comments?**

1-800-719-9260

**Consumer Information****What you should know about Ranitidine Tablets, USP 75 mg Acid Reducer****Ranitidine Tablets, USP 75 mg**

(Please read all of this information before taking Ranitidine Tablets, USP 75 mg. Save this leaflet for future reference.)

**What is Ranitidine Tablet 75 mg?**

- Ranitidine Tablet contains 75 mg of ranitidine (as ranitidine hydrochloride, 84 mg), a medicine that doctors have prescribed more than 200 million times worldwide.


**Excellent safety record**

- Ranitidine Tablets 75 mg has been used safely and effectively for years as an over-the-counter medication.
- Ranitidine Tablets 75 mg are sugar free.

**What symptoms does Ranitidine Tablet 75 mg treat and prevent?**

Ranitidine Tablet 75 mg relieves and prevents heartburn associated with acid indigestion and sour stomach.

Certain foods or beverages, and even lying down to sleep, can cause heartburn associated with acid indigestion or sour stomach. It is normal for the stomach to produce acid, especially after consuming food or beverages. However, acid in the wrong place, such as the esophagus, or too much acid, can cause burning pain and discomfort.



### How should I take Ranitidine Tablet 75 mg?

- To **relieve** symptoms, swallow 1 tablet with a glass of water.
- To **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn.

This medicine can be used up to twice daily (do not take more than 2 tablets in 24 hours). Ranitidine Tablets 75 mg should not be given to children under 12 years old unless directed by a doctor.

### How does Ranitidine Tablet 75 mg work?

Ranitidine Tablet 75 mg reduces the production of stomach acid. This is what makes Ranitidine Tablets 75 mg different from antacids, which neutralize the acid already in your stomach. Antacids do not reduce the production of acid.

### Additional tips for managing heartburn

- Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables
- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

### When should I see a doctor?

- **Allergy alert:** Do not use if you are allergic to ranitidine or other acid reducers.
- **Do not use** if you have

- trouble or pain swallowing food
- vomiting with blood
- bloody or black stools

These may be signs of a serious condition. See your doctor.

- with other acid reducers
- **Ask a doctor before use if you have**
- had heartburn over 3 months.

This may be a sign of a more serious condition.

- heartburn with **lightheadedness,**
- **sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- **Stop use and ask a doctor if**
- your heartburn continues or worsens
- you need to take this product for more than 14 days
- **If pregnant or breast-feeding,** ask a health professional before use.
- **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Questions?** Call 1-800-719-9260

**BOTTLES: Bottle is sealed with printed foil under cap. Do not use if foil is open or torn. Bottle contains desiccant.**

**BLISTERS: Do not use if the individual blister unit is open or torn.**

**DISTRIBUTED BY PERRIGO, ALLEGAN, MI 49010**

**Made in India**

:27100 00 J2

### **Principal Display Panel**

Ranitidine Tablets 75 mg

ACID REDUCER

Compare to the active ingredient of Zantac 75®

Prevents and Relieves

Heartburn Associated with Acid Indigestion & Sour Stomach

# DOSES {Replace "#" with number of tablets in package }

EXP.  
: 27J39 CZ CI

LOT NO.



exchange select 

# Ranitidine Tablets 75 mg ACID REDUCER

Compare to the active ingredient of Zantac 75®

**Prevents and Relieves**  
Heartburn Associated with  
Acid Indigestion & Sour Stomach

**30 TABLETS**  
**(30 DOSES)**

exchange select 

Ranitidine Tablets 75 mg  
**ACID REDUCER**  
Prevents and Relieves  
Heartburn Associated with Acid Indigestion & Sour Stomach



**Drug Facts (continued)**

**Other information**

- do not use if printed foil under cap is broken or missing
- store at 20° - 25°C (68° - 77°F)
- avoid excessive heat or humidity
- this product is sugar free

**Inactive ingredients** colloidal silicon dioxide, croscarmellose sodium, diethyl phthalate, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, titanium dioxide.

**Questions or comments? 1-800-719-9260**

Bottle contains desiccant.

**SATISFACTION GUARANTEED**

OR YOUR MONEY BACK

Made in India  
Packed For: Your Military Exchanges  
By: Perrigo Company, Allegan, MI USA 49010



IMPORTANT: Read the directions, consumer information leaflet and warnings before use. Keep the carton. It contains important information.

**Drug Facts**

**Active ingredient (in each tablet)**

Ranitidine 75 mg (as ranitidine hydrochloride 84 mg).....Acid reducer

**Purpose**

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

**Warnings**

**Allergy alert:** Do not use if you are allergic to ranitidine or other acid reducers

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

**Ask a doctor before use if you have**

- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- adults and children 12 years and over:
  - to relieve symptoms, swallow 1 tablet with a glass of water
  - to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn
  - can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

**EXCHANGE SELECT ACID REDUCER**

ranitidine tablet

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55301-271
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
RANITIDINE HYDROCHLORIDE (UNII: BK76465IHM) (RANITIDINE - UNII:884KT10YB7)	RANITIDINE	75 mg

**Inactive Ingredients**

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
DIETHYL PHTHALATE (UNII: UF064M00AF)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	

<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	PINK	<b>Score</b>	no score
<b>Shape</b>	HEXAGON (6 sided)	<b>Size</b>	8mm
<b>Flavor</b>		<b>Imprint Code</b>	W;75
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55301-271-39	1 in 1 CARTON		
1		30 in 1 BOTTLE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076760	09/12/2012	

**Labeler** - Army & Air Force Exchange Service (001695568)

Revised: 10/2012

Army & Air Force Exchange Service