

**READY AMERICA FIRST AID OUTDOOR- 107 PIECES- aspirin, ibuprofen, acetaminophen, alcohol, benzalkonium chloride, bacitracin zinc, neomycin sulfate, polymyxin b sulfate, benzalkonium chloride, lidocaine, benzocaine, alcohol  
Genuine First Aid, LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Ready America First Aid Outdoor Kit 107 pcs**

Active Ingredient: .....Bacitracin Zinc 400 units

Neomycin Sulfate 5mg ( equivalent to 3.5 mg Neomycin base)

Polymyxin B Sulfate 5000 units

Uses: To help prevent infection in:  
minor cuts; scrapes; burns

Warnings:

For external use only.

Do not use: in eyes; over large areas of the body;

If allergic to any of the ingredients; for more than one week unless directed by a physician.

Stop use and consult a doctor:

if the condition persists or gets worse; a rash or other allergic reaction develops

Keep out of reach of children.

If ingested, contact a Poison

Control Center right away.

Directions: clean affected area; apply small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily; may be covered with a sterile bandage

Other information:

Store at room temperature.

Genuine Triple Antibiotic

First Aid Ointment

To Help Prevent Infection

Each Gram Contains:

Bacitracin Zinc 400 units

Neomycin Sulfate 5 mg

(equivalent to 3.5 mg

Neomycin base)

Polymyxin B Sulfate 5000 units

Net Wt. 0.5g ; (1/64 oz)

Manufactured in CHINA for

GENUINE FIRST AID.

| Active ingredient (in each tablet)     | Purpose                     |
|--|-----------------------------|
| Ibuprofen USP (NSAID*) 200mg . . . . . | Pain reliever/fever reducer |
| *nonsteroidal anti-inflammatory drug   |                             |

Uses temporarily relieves minor aches and pains due to:

the common cold

headache

toothache

muscular aches

backache

minor pain of arthritis

menstrual cramps temporarily reduces fever

### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: shock, facial swelling, asthma (wheezing) rash, skin reddening, blisters, hives. If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinner (anticoagulant) or steroid drug, take other drugs containing NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer, right before or after heart surgery.

Ask a doctor before use if stomach bleeding warning applies to you; you have a history of stomach problems such as heartburn; you have a high blood pressure, heart disease, liver cirrhosis, or kidney disease; you are taking a diuretic

Ask a doctor before use if you are: taking any other drug containing NSAID (prescription or nonprescription); taking aspirin for heart attack or stroke, because Ibuprofen may decrease this benefit of aspirin; taking any other drug

When using this product:  
take with food or milk if stomach upset occurs

Stop use and ask a doctor If:  
you experience any of the following signs of stomach bleeding; feel faint; vomit blood; have bloody or

black stools; have stomach pain that does get better; pain gets worse or lasts more than 10 days; fever gets worse or lasts more than 3 days; redness or swelling is present in the painful area; any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions:**

do not use more than directed; the smallest effective dose should be used; do not take longer than 10 days, unless directed by a doctor.

Adults and Children (12 years and older): Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

Other information: Store at controlled room temperature; avoid excessive heat 40 degree Celsius (104 degree Fahrenheit); tamper evident sealed packets; do not use any opened or torn packets

Inactive ingredients: cellulose, corn starch, fumed silica gel, hypromellose, lactose, magnesium stearate, polydextrose, polyethylene glycol, povidone, silica, sodium starch glycolate, stearic acid, titanium dioxide, triacetin.

Distributed by GENUINE FIRST AID  
600 Cleveland Str Suite 400, Clearwater, FL 33755

IBUPROFEN 2 Tablets

IBUPROFEN 2 Tablets

| Active Ingredient (in each tablet) | Purpose                       |
|------------------------------------|-------------------------------|
| Aspirin (NSAID*) 325 mg.....       | Pain Reliever / fever reducer |

\*nonsteroidal anti-inflammatory drug

Uses Temporarily relieves minor aches and pains associated with:  
headache ; muscular aches ; minor arthritis pain ; backache ; common cold ; toothache ; menstrual cramps ; Temporarily reduces fever

**Warnings**

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: hives, skin reddening, facial swelling, rash, asthma (wheezing), blisters, shock, If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
are age 60 or older; have had stomach ulcers or bleeding problems; take a blood thinner (anticoagulant)

or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others); have 3 or more alcoholic drinks every day while using this product; take more or for a longer time than directed

Do not use: if you have ever had an allergic reaction to any other pain reliever/ fever reducer; right before or after heart surgery; if you are taking a prescription drug for gout, diabetes or arthritis

Ask a doctor before use if: stomach bleeding warning applies to you; you have a history of stomach problems such as heartburn; you have high blood pressure, heart disease, liver cirrhosis, or kidney disease; you are taking a diuretic

Ask a doctor or pharmacist before use if you are:

under a doctor's care for any serious condition; taking any other drug

When using this product: take with food or milk if stomach upset occurs

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding:

feel faint; vomit blood; have bloody or black stools; have stomach

pain that does not get better; pain gets worse or lasts more than 10 days; fever gets worse or lasts more than 3 days; you have difficulty swallowing; if ringing in the ears or loss of hearing occurs; redness or swelling is present in the painful areas; any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

#### Directions

do not use more than directed

the smallest effective dose should be used

drink a full glass of water with each dose

do not take longer than 10 days, unless directed by a doctor

Adults and children: (12 years and older) Take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

Store at 59 - 86 degree Fahrenheit (15 - 30 degree Celsius); avoid excessive heat and humidity; tamper evident sealed packets;

Do not use any opened or torn packets

Inactive Ingredients: hypromellose, polyethylene glycol, propylene glycol, corn starch

Distributed by GENUINE FIRST AID

600 Cleveland Str Suite 400, Clearwater, FL 33755

ASPIRIN 2 Tablets

Active Ingredient:

Purpose

Benzalkonium Chloride 0.40%..... First Aid Antiseptic

Use: For Professional and Hospital use. Helps prevent infection. Antiseptic cleansing of face, hands and body without soap and water.

Warning: For external use only.

Keep out of reach of children: If swallowed, get medical help or contact a Poison Control Center right away.

Stop use if unusual redness, swelling or other symptoms occur. Consult a physician immediately.

Do not use in the eyes or over large areas of the body.

Directions: Tear open packet, unfold towelette and use to cleanse desired skin area. Discard towelette appropriately after single use.

Inactive Ingredient: Purified water

LOT/EXP:                      Made in CHINA

20130301

Antiseptic Towelette

Genuine First Aid LLC, Clearwater FL 33755

[www.GenuineFirstAid.com](http://www.GenuineFirstAid.com)

1/pouch

GENUINE FIRST AID

### **Active Ingredient (in each tablet)**

Acetaminophen 325 mg

### **Purpose**

Analgesic/antipyretic

### **Uses**

temporary relief of minor aches and pains associated with:

common cold; headache; toothache; muscular aches; backache; arthritis; menstrual cramps; and reduction of fever

### **Warnings**

Liver warning: This product contains acetaminophen.

Severe liver damage may occur if: adult takes more than 12 tablets in 24 hours, which is the maximum daily amount; child takes more than 5 doses in 24 hours; taken with other drugs containing acetaminophen; adult has 3 or more alcoholic drinks every day while using this product

### **Do not use**

with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist; for more than 10 days for pain unless directed by a doctor; for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if the user has liver disease

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

Stop use and ask a doctor if: symptoms do not improve; pain gets worse or lasts for more than 10 days; fever gets worse or lasts for more than 3 days; new symptoms occur; redness or swelling is present; a rare sensitivity reaction occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, contact a doctor or Poison Control Center immediately. Prompt

medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Do not exceed recommended dosage

### **Inactive Ingredients**

Cornstarch, polyethylene glycol, stearic acid, povidone

#### Directions

|  |  |
|--|--|
| Adults and Children<br>12 years of age<br>or older | Take 2 tablets every 4 to 6 hours as<br>needed. Do not take more than 12 tablets<br>in 24 hours. |
|--|--|

|                               |  |
|-------------------------------|--|
| Children 6-11 years<br>of age | Take 1 tablet every 4 to 6 hours as<br>needed. Do not take more than 5<br>tablets in 24 hours. |
|-------------------------------|--|

|                                  |   |
|----------------------------------|---|
| Children under 6<br>years of age | Do not use this regular strength product.<br>This will provide more than the<br>recommended dose (overdose) and could<br>cause serious health problems. |
|----------------------------------|---|

Store at 59-86 degree F (15-30 degree C)

tamper evident sealed packets; do not use any open or torn packets

Distributed by GENUINE FIRST AID  
600 Cleveland Str Suite 400, Clearwater, FL 33755

GENUINE FIRST AID      2 Tablets

NON-ASPIRIN

### **Active Ingredient**

Active Ingredient:

Isopropyl Alcohol, 70% v/v

### **Purpose**

Antiseptic

### **Use**

For preparation of skin before injection.

### **Warnings**

For external use only.

Flammable - keep away from fire or flame

Store at room temperature 15-30 degree Celsius (59-86 degree Fahrenheit)

Do not use: with electrocautery, in the eyes.

Stop use if irritation and redness develop. If condition persists for more than 72 hours, consult your doctor.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

### **Directions**

Wipe Injection site vigorously and discard.

### **Inactive Ingredient**

Purified water.

LOT/EXP:        Made in CHINA

20140301

Alcohol Cleansing Pad  
Genuine First Aid LLC, Clearwater FL 33755

[www.GenuineFirstAid.com](http://www.GenuineFirstAid.com)

1/pouch

GENUINE FIRST AID

100 Alcohol cleansing pads

Reorder GFAP-39-02

### **Active Ingredients**

ACTIVE INGREDIENTS:

Benzalkonium Chloride 0.13%  
Lidocaine HCL 0.5%

## **Purpose**

Purpose: First aid antiseptic, external analgesic

## **Uses**

First aid to help prevent infection and for the temporary relief of pain and itching associated with:

Minor Cuts

Scrapes

Burns

## **Warnings**

For external use only

Do not use: In eyes, in large quantities, over raw blistered areas, or on deep puncture wounds, animal bites or serious burns, for more than one week

Do not use:

in the eyes or apply over large areas of the body.

longer than 1 week unless directed by a doctor.

in large quantities, particularly over raw surfaces or blistered areas.

Ask a doctor before use if you have deep puncture wounds, animal bites or serious burns.

When using this product, avoid contact with the eyes.

Stop use and ask a doctor if  
condition worsens

symptoms persist for more than 7 days

condition clears up and occurs again within a few days

Keep out of reach of Children.

If ingested, contact a Poison Control Center right away.

## **Directions**

Adults and children 2 years of age and older

clean affected area.



apply a small amount of this product on the area 1 to 3 times daily.

may be covered with a sterile bandage

children under 2 years of age: consult a doctor

Other Information:

Store at room temperature (do not freeze).

Taper evident sealed packets.

Do not use packet if opened or torn.

### **Inactive Ingredients**

Aloe vera, cetyl alcohol, diazolidinyl urea, edetate disodium, glycerin, glyceryl monostearate, methylparaben, mineral oil, polyethylene glycol, propylene glycol, propylparaben, purified water, stearic acid, irolamine

LOT/EXP:

Made in CHINA

20130301

GFA Production Xiamen Co., Ltd

No. 20 Huli Industrial Park, Meixi Road, Tong'an, Xiamen, Fujian, China 361100

Tel: 86-592-7269515 Fax: 86-592-7269528 Http: //www.gfaproduction.com

Genuine First Aid Burn Cream

Antiseptic Pain Relief With Aloe

Net Wt 0.9g (1/32 oz)

Manufactured in CHINA for

Genuine First Aid.

### **Active Ingredient**

Active Ingredient:

Purpose:

Benzocaine, 6% w/v

SD alcohol, 60% w/v

### **Purpose**

Topical Anesthetic

Antiseptic

### **Uses**

Use: For the temporary relief of pain and itching associated with minor burns, scrapes and insect bites.

**Directions**

Clean intended area thoroughly with pad. Discard after single use.

**Warnings**

Warnings: For external use only.

Avoid contact with eyes. If this happens, rinse thoroughly with water.

**Keep out of reach of children**

If swallowed, get medical help or contact a Poison Control Center right away.

**Inactive Ingredients**

Purified water

Flammable - keep away from fire or flame.

Do not use: In eyes, on broken skin, deep puncture wounds. If unusual redness, swelling, irritation or other symptoms occur, consult a physician immediately.

Made in CHINA

LOT/EXP:

Insect Sting Relief Pad

Genuine First Aid LLC, Clearwater FL 33755

[www.GenuineFirstAid.com](http://www.GenuineFirstAid.com)

1/pouch

GENUINE FIRST AID

REORDER AST-001

**Antiseptic  
Towelette**  
**Toallitas  
Antisepicas**



Genuine First Aid LLC, Clearwater FL 33755  
www.GenuineFirstAid.com



**GENUINE FIRST AID®**



DRUG FACTS - Antiseptic Towelette

Active Ingredient: Benzalkonium Chloride 0.40% Purpose: First Aid Antiseptic

Use: For Professional and Hospital use. Helps prevent infection. Antiseptic cleansing of face, hands and body without soap and water.

Warnings: For external use only. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. If unusual redness, swelling or other symptoms occur, consult a physician immediately.

Do not use: In the eyes, or over large areas of the body.

Directions: Tear open packet, unfold towelette and use to cleanse desired skin area. Discard towelette appropriately after single use.

Inactive ingredient: Purified water

Made in CHINA

LOT/EXP:  
20130301

--- TEAR HERE ---

Reorder TAO-001

# Genuine Triple Antibiotic

**First Aid Ointment  
To Help Prevent Infection**

**Each Gram Contains:**  
**Bacitracin Zinc 400 units**  
**Neomycin Sulfate 5 mg**  
**(equivalent to 3.5 mg**  
**Neomycin base)**  
**Polymyxin B Sulfate 5000 units**  
**Net Wt 0.5g • (1/64 oz)**



Manufactured in CHINA for  
**GENUINE FIRST AID.**

## Drug Facts

Uses: To help prevent infection  
in • minor cuts • scrapes • burns

## Warnings:

For external use only

Do not use • in eyes • over large  
areas of the body • if allergic to  
any of the ingredients • for more  
than one week unless directed  
by a physician

Stop use and consult a doctor  
• if the condition persists or gets  
worse • a rash or other allergic  
reaction develops

Keep out of reach of children.

If ingested, contact a Poison  
Control Center right away.

Directions: • clean affected area  
• apply small amount of this product  
(an amount equal to the surface  
area of the tip of a finger) on the  
area 1 to 3 times daily • may be  
covered with a sterile bandage

Other information:

Store at room temperature

LOT/EXP:

20130118

## IBUPROFEN 2 Tablets

**Active ingredient (in each tablet)** Purpose  
Ibuprofen USP (NSAID)\* 200mg . . . . . Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses** temporarily relieves minor aches and pains due to:  
■ the common cold ■ headache ■ toothache ■ muscular aches  
■ backache ■ minor pain of arthritis ■ menstrual cramps  
temporarily reduces fever

**Warnings** **Allergy alert:** Ibuprofen may cause a severe  
allergic reaction, especially in people allergic to aspirin. Symptoms  
may include: ■ shock ■ facial swelling ■ asthma (wheezing)  
■ rash ■ skin reddening ■ blisters ■ hives If an allergic reaction  
occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains a nonsteroidal  
anti-inflammatory drug (NSAID), which may cause severe stomach  
bleeding. The chance is higher if you: ■ are age 60 or older  
■ have had stomach ulcers or bleeding problems ■ take a blood  
thinner (anticoagulant) or steroid drug ■ take other drugs containing  
NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more  
alcoholic drinks every day while using this product ■ take more or  
for a longer time than directed

**Do not use** ■ if you have ever had an allergic reaction to any other  
pain reliever/fever reducer ■ right before or after heart surgery

**Ask a doctor before use if** ■ stomach bleeding warning applies to  
you ■ you have a history of stomach problems such as heartburn  
■ you have high blood pressure, heart disease, liver cirrhosis, or  
kidney disease ■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are** ■ taking any  
other drug containing an NSAID (prescription or nonprescription) ▶

## Warnings (continued)

■ taking aspirin for heart attack or stroke, because ibuprofen may  
decrease this benefit of aspirin ■ taking any other drug

## When using this product

■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if** ■ you experience any of the  
following signs of stomach bleeding: ■ feel faint ■ vomit blood  
■ have bloody or black stools ■ have stomach pain that does not  
get better ■ pain gets worse or lasts more than 10 days ■ fever  
gets worse or lasts more than 3 days ■ redness or swelling is present  
in the painful area ■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It  
is especially important not to use ibuprofen during the last 3 months of  
pregnancy unless definitely directed to do so by a doctor because it may  
cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical  
help or contact a Poison Control Center right away.

**Directions** ■ do not use more than directed ■ the smallest  
effective dose should be used ■ do not take longer than 10 days,  
unless directed by a doctor

**Adults and Children (12 years and older):** Take 1 tablet every 4 to 6  
hours while symptoms persist. If pain or fever does not respond to 1  
tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours,  
unless directed by a doctor.

**Children under 12 years:** Do not give to children under 12 years of age.

**Other information** ■ store at controlled room temperature ■ avoid  
excessive heat 40° C (104° F) ■ tamper evident sealed packets ■ do not  
use any opened or torn packets

**Inactive ingredients** cellulose, corn starch, fumed silica gel,  
hypromellose, lactose, magnesium stearate, polydextrose, polyethylene  
glycol, povidone, silica, sodium starch glycolate, stearic acid, titanium  
dioxide, triacetin

Distributed by GENUINE FIRST AID  
600 Cleveland Str Suite 400, Clearwater, FL 33755

# ASPIRIN

2 Tablets

---

**Active ingredient (in each tablet)****Purpose**

Aspirin (NSAID\*) 325 mg . . . . . Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

---

**Uses** Temporarily relieves minor aches and pains associated with

- headache ■ muscular aches ■ minor arthritis pain ■ backache
- common cold ■ toothache ■ menstrual cramps

Temporarily reduces fever

---

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ skin reddening ■ facial swelling ■ rash ■ asthma (wheezing) ■ blisters ■ shock  
If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
■ are age 60 or older ■ have had stomach ulcers or bleeding problems  
■ take a blood thinner (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery ■ if you are taking a prescription drug for gout, diabetes or arthritis

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ▶

---

---

**Warnings (continued)**

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition ■ taking any other drug

**When using this product** ■ take with food or milk if stomach upset occurs

---

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:  
■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days ■ you have difficulty swallowing
  - if ringing in the ears or loss of hearing occurs ■ redness or swelling is present in the painful area ■ any new symptoms appear
- 

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

---

**Directions**

- do not use more than directed
  - the smallest effective dose should be used
  - drink a full glass of water with each dose
  - do not take longer than 10 days, unless directed by a doctor
- 

**Adults and children: (12 years and older)** Take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.

**Children under 12 years:** Do not give to children under 12 years of age.

---

**Other information**

- store at 59°-86°F (15°-30°C) ■ avoid excessive heat and humidity ■ tamper evident sealed packets ■ do not use any opened or torn packets
- 

**Inactive ingredients** hypromellose, polyethylene glycol, propylene glycol, corn starch

---

Distributed by GENUINE FIRST AID  
600 Cleveland Str Suite 400, Clearwater, FL 33755

REORDER ACP-001

**Alcohol Cleansing  
Pad**

**Toallitas Humedas  
con Alcohol**



ISO 9001:2010



Genuine First Aid LLC, Clearwater FL 33755  
www.GenuineFirstAid.com



**GENUINE FIRST AID®**



**DRUG FACTS - Alcohol Cleansing Pad**

**Active Ingredient:** Isopropyl Alcohol, 70% v/v . . . . **Purpose:** Antiseptic

**Use:** For preparation of the skin before injection.

**Warnings: For external use only.**

**Flammable - keep away from fire or flame.**

**Do not use:** with electrocautery, in the eyes

**Stop use** if irritation and redness develop.

If condition persists for more than 72 hours,  
consult your doctor.

**Keep out of reach of children.** If swallowed,  
get medical help or contact a Poison Control  
Center right away.

**Directions:** Wipe injection site vigorously and  
discard.

**Other information:** Store at room temperature  
15°-30° C (59°-86° F)

**Inactive ingredient:** Purified water.

----- TEAR HERE -----

LOT/EXP:

Made in CHINA

20130301



GENUINE FIRST AID. 2 Tablets

# NON-ASPIRIN

**Active ingredient (in each tablet) Purpose**  
Acetaminophen 325 mg . . . . . Analgesic/antipyretic

### Uses

temporary relief of minor aches and pains associated with

- common cold    ■ headache    ■ toothache
- muscular aches    ■ backache    ■ arthritis
- menstrual cramps    ■ and reduction of fever

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if ■ adult takes more than 12 tablets in 24 hours, which is the maximum daily amount ■ child takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist ■ for more than 10 days for pain unless directed by a doctor ■ for more than 3 days for fever unless directed by a doctor

**Ask a doctor before use if the user has liver disease**

**Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin** ▶

### Warnings (continued)

**Stop use and ask a doctor if** ■ symptoms do not improve ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ new symptoms occur ■ redness or swelling is present ■ a rare sensitivity reaction occurs

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of accidental overdose, contact a doctor or Poison Control Center immediately. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. **Do not exceed recommended dosage.**

### Directions

|   |  |
|---|--|
| Adults and children 12 years of age and older | Take 2 tablets every 4 to 6 hours as needed. Do not take more than 12 tablets in 24 hours.   |
| Children 6-11 years of age                    | Take 1 tablet every 4 to 6 hours as needed. Do not take more than 5 tablets in 24 hours.   |
| Children under 6 years of age                 | Do not use this regular strength product. This will provide more than the recommended dose (overdose) and could cause serious health problems. |

**Other information** ■ store at 59°-86°F (15°-30°C) ■ tamper evident sealed packets ■ do not use any open or torn packets

**Inactive ingredients** corn starch, hydroxypropyl methylcellulose, polyethylene glycol, pregelatinized starch, stearic acid. May contain povidone and sodium starch glycolates.

Distributed by GENUINE FIRST AID  
600 Cleveland Str Suite 400, Clearwater, FL 33755

Reorder BC-001

# Genuine First Aid Burn Cream

**Antiseptic Pain Relief  
With Aloe**

## ACTIVE INGREDIENTS:

**Benzalkonium  
Chloride 0.13%  
Lidocaine HCL 0.5%**

**Net Wt 0.9g • (1/32 oz)**



Manufactured in CHINA for  
**GENUINE FIRST AID.**

## Drug Facts

**Purpose:** First aid antiseptic,  
external analgesic

**Uses:** First aid to help prevent  
infection and for the temporary  
relief of pain and itching  
associated with: • minor cuts  
• scrapes • burns

## Warnings:

For external use only

Do not use • in eyes • in large  
quantities • over raw blistered  
areas, or on deep puncture  
wounds, animal bites or serious  
burns • for more than one week  
Keep out of reach of children.

If ingested, contact a Poison  
Control Center right away.

**Directions:** • clean affected area  
• apply small amount not more  
than 3 times daily • may be  
covered with a sterile bandage

## Other information:

Store at room temperature

**LOT/EXP:**

20130301

REORDER GFAP-29-07

# Insect Sting Relief Pad

---

## Toallitas para Picaduras de Insectos



GFA Production Xiamen Co., Ltd  
[www.gfaproduction.com](http://www.gfaproduction.com)



 **GFA Production Xiamen Co., Ltd**  
No. 20 Huli Industrial Park, Meidi Road,  
Tongjiao, Xiamen, Fujian, China 361100  
 **Wellkang Ltd t/a Wellkang Tech Consulting**  
Suite B, 20 Harley Street  
LONDON W1G 9QR, England, United Kingdom



## DRUG FACTS - Insect Sting Relief Pad

**Active Ingredient:** Benzocaine, 6% w/v ..... **Purpose:** Topical Anesthetic  
SD alcohol, 60% w/v ..... Antiseptic

**Use:** For the temporary relief of pain and itching  
associated with minor burns, scrapes and insect  
bites.

**Warnings:** For external use only.

**Do not use:** In eyes, on broken skin, deep  
puncture wounds. If unusual redness, swelling,  
irritation or other symptoms occur, consult a  
physician immediately.

**Inactive Ingredients:** Purified water.

**DIRECTIONS:** clean intended area thoroughly with pad.  
Discard after single use.

**Keep out of reach of children.** If swallowed, get medical  
help or contact a Poison Control Center right away.

Flammable - keep away from fire or flame. Avoid contact  
with eyes. If this happens, rinse thoroughly with water.

**LOT/EXP:**

20140115

----- TEAR HERE -----



This kit contains dated items. Please check before use.

20130103  
 20160103  
 20130103



www.readyamerica.com  
 Escondido, CA 92029  
 for Ready America, Inc.  
 Manufactured in China



**107**  
 pieces  
 107 piezas

**First Aid Outdoor Kit**

Kit de primeros auxilios para fuera de casa



Kit de primeros auxilios para fuera de casa

**First Aid Outdoor Kit**



• **107 pieces**

• Includes **critical supplies** (for wound care) not found in other kits

• **Be prepared** away from home

• Recommended by **Health Care Professionals**



• **107 piezas**

• Incluye artículos críticos para el cuidado de heridas que no se encuentran en otros equipos

• Estar preparados fuera de casa

• Recomendado por profesionales de la salud



**READY AMERICA FIRST AID OUTDOOR- 107 PIECES**

aspirin, ibuprofen, acetaminophen, alcohol, benzalkonium chloride, bacitracin zinc, neomycin sulfate, polymyxin b sulfate, benzalkonium chloride, lidocaine, benzocaine, alcohol kit

**Product Information**

|              |                |                    |                |
|--------------|----------------|--------------------|----------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:52124-4002 |
|--------------|----------------|--------------------|----------------|

**Packaging**

| # | Item Code        | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------|----------------------|--------------------|
| 1 | NDC:52124-4002-8 | 1 in 1 KIT          |                      |                    |

## Quantity of Parts

| Part # | Package Quantity | Total Product Quantity |
|--------|------------------|------------------------|
| Part 1 | 1 TUBE           | 0.5 g                  |
| Part 2 | 1 PACKET         | 2                      |
| Part 3 | 1 PACKAGE        | 2                      |
| Part 4 | 3 PACKAGE        | 1.5 mL                 |
| Part 5 | 1 PACKAGE        | 2                      |
| Part 6 | 3 PACKAGE        | 2.4 mL                 |
| Part 7 | 2 PACKET         | 1.8 g                  |
| Part 8 | 2 PACKAGE        | 1 mL                   |

## Part 1 of 8

### GENUINE TRIPLE ANTIBIOTIC

bacitracin zinc,neomycin sulfate,polymyxin b sulfate ointment

## Product Information

|                         |                |
|-------------------------|----------------|
| Item Code (Source)      | NDC:52124-0003 |
| Route of Administration | TOPICAL        |

## Active Ingredient/Active Moiety

| Ingredient Name  | Basis of Strength | Strength         |
|--|-------------------|------------------|
| BACITRACIN ZINC (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)      | BACITRACIN        | 400 [iU] in 1 g  |
| NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)       | NEOMYCIN          | 5 mg in 1 g      |
| POLYMYXIN B SULFATE (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K) | POLYMYXIN B       | 5000 [iU] in 1 g |

## Inactive Ingredients

| Ingredient Name          | Strength |
|--------------------------|----------|
| WATER (UNII: 059QF0KO0R) |          |

## Packaging

| # | Item Code        | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------|----------------------|--------------------|
| 1 | NDC:52124-0003-1 | .5 g in 1 TUBE      |                      |                    |

## Marketing Information

| Marketing Category  | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part333B                                 | 01/09/2013           |                    |

## Part 2 of 8

# IBUPROFEN

ibuprofen tablet

## Product Information

|                         |                |
|-------------------------|----------------|
| Item Code (Source)      | NDC:52124-0009 |
| Route of Administration | ORAL           |

## Active Ingredient/Active Moiety

| Ingredient Name  | Basis of Strength | Strength |
|--|-------------------|----------|
| IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM) | IBUPROFEN         | 200 mg   |

## Inactive Ingredients

| Ingredient Name                         | Strength |
|---|----------|
| POWDERED CELLULOSE (UNII: SMD1X3XO9M)   |          |
| STARCH, CORN (UNII: O8232NY3SJ)         |          |
| HYPROMELLOSES (UNII: 3NXW29V3WO)        |          |
| LACTOSE (UNII: J2B2A4N98G)              |          |
| MAGNESIUM STEARATE (UNII: 70097M6I30)   |          |
| POLYDEXTROSE (UNII: VH2XOU12IE)         |          |
| POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A) |          |
| POVIDONE (UNII: FZ989GH94E)             |          |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4)      |          |
| STEARIC ACID (UNII: 4ELV7Z65AP)         |          |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP)     |          |
| TRIACETIN (UNII: XHX3C3X673)            |          |

## Product Characteristics

|          |               |              |          |
|----------|---------------|--------------|----------|
| Color    | white (WHITE) | Score        | no score |
| Shape    | ROUND         | Size         | 10mm     |
| Flavor   |               | Imprint Code | 44;352   |
| Contains |               |              |          |

## Packaging

| # | Item Code        | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------|----------------------|--------------------|
| 1 | NDC:52124-0009-1 | 2 in 1 PACKET       |                      |                    |

## Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA               | ANDA075010                               | 01/09/2013           |                    |

## Part 3 of 8

### NON-ASPIRIN

acetaminophen tablet

#### Product Information

Item Code (Source) NDC:52124-0010

Route of Administration ORAL

#### Active Ingredient/Active Moiety

| Ingredient Name  | Basis of Strength | Strength |
|--|-------------------|----------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN     | 325 mg   |

#### Inactive Ingredients

| Ingredient Name                         | Strength |
|---|----------|
| STARCH, CORN (UNII: O8232NY3SJ)         |          |
| POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A) |          |
| STEARIC ACID (UNII: 4ELV7Z65AP)         |          |
| POVIDONE (UNII: FZ989GH94E)             |          |

#### Product Characteristics

|          |               |              |          |
|----------|---------------|--------------|----------|
| Color    | white (White) | Score        | no score |
| Shape    | ROUND         | Size         | 11mm     |
| Flavor   |               | Imprint Code | AZ;234   |
| Contains |               |              |          |

#### Packaging

| # | Item Code        | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------|----------------------|--------------------|
| 1 | NDC:52124-0010-1 | 2 in 1 PACKAGE      |                      |                    |

#### Marketing Information

| Marketing Category      | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|-------------------------|--|----------------------|--------------------|
| OTC monograph not final | part343                                  | 01/09/2013           |                    |

## Part 4 of 8

### ALCOHOL CLEANSING PAD

isopropyl alcohol liquid

**Product Information**

|                           |                |
|---------------------------|----------------|
| <b>Item Code (Source)</b> | NDC:52124-0002 |
|---------------------------|----------------|

|                                |         |
|--------------------------------|---------|
| <b>Route of Administration</b> | TOPICAL |
|--------------------------------|---------|

**Active Ingredient/Active Moiety**

| <b>Ingredient Name</b>   | <b>Basis of Strength</b> | <b>Strength</b>    |
|--|--------------------------|--------------------|
| ISOPROPYL ALCOHOL (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302) | ISOPROPYL ALCOHOL        | 70 mL<br>in 100 mL |

**Inactive Ingredients**

| <b>Ingredient Name</b>   | <b>Strength</b> |
|--------------------------|-----------------|
| WATER (UNII: 059QF0K00R) |                 |

**Packaging**

| # | <b>Item Code</b> | <b>Package Description</b> | <b>Marketing Start Date</b> | <b>Marketing End Date</b> |
|---|------------------|----------------------------|-----------------------------|---------------------------|
| 1 | NDC:52124-0002-1 | 0.5 mL in 1 PACKAGE        |                             |                           |

**Marketing Information**

| <b>Marketing Category</b> | <b>Application Number or Monograph Citation</b> | <b>Marketing Start Date</b> | <b>Marketing End Date</b> |
|---------------------------|---|-----------------------------|---------------------------|
| OTC monograph not final   | part333A  | 01/09/2013                  |                           |

**Part 5 of 8****ASPIRIN**

aspirin tablet

**Product Information**

|                           |                |
|---------------------------|----------------|
| <b>Item Code (Source)</b> | NDC:52124-0011 |
|---------------------------|----------------|

|                                |      |
|--------------------------------|------|
| <b>Route of Administration</b> | ORAL |
|--------------------------------|------|

**Active Ingredient/Active Moiety**

| <b>Ingredient Name</b>                                 | <b>Basis of Strength</b> | <b>Strength</b> |
|--|--------------------------|-----------------|
| ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E) | ASPIRIN                  | 325 mg          |

**Inactive Ingredients**

| <b>Ingredient Name</b> | <b>Strength</b> |
|------------------------|-----------------|
|------------------------|-----------------|

|   |  |
|---|--|
| HYPROMELLOSES (UNII: 3NXW29V3WO)        |  |
| POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A) |  |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3)     |  |
| STARCH, CORN (UNII: O8232NY3SJ)         |  |

### Product Characteristics

|          |               |              |                |
|----------|---------------|--------------|----------------|
| Color    | white (white) | Score        | no score       |
| Shape    | ROUND         | Size         | 11mm           |
| Flavor   |               | Imprint Code | 44;157;ASPIRIN |
| Contains |               |              |                |

### Packaging

| # | Item Code        | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------|----------------------|--------------------|
| 1 | NDC:52124-0011-1 | 2 in 1 PACKAGE      |                      |                    |

### Marketing Information

| Marketing Category  | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part343                                  | 01/09/2013           |                    |

### Part 6 of 8

#### ANTISEPTIC

benzalkonium chloride swab

### Product Information

|                         |                |
|-------------------------|----------------|
| Item Code (Source)      | NDC:52124-0001 |
| Route of Administration | TOPICAL        |

### Active Ingredient/Active Moiety

| Ingredient Name   | Basis of Strength     | Strength       |
|---|-----------------------|----------------|
| BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y) | BENZALKONIUM CHLORIDE | 1.3 mg in 1 mL |

### Inactive Ingredients

| Ingredient Name          | Strength |
|--------------------------|----------|
| WATER (UNII: 059QF0KO0R) |          |

### Packaging

| # | Item Code        | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------|----------------------|--------------------|
| 1 | NDC:52124-0001-1 | 0.8 mL in 1 PACKAGE |                      |                    |

## Marketing Information

| Marketing Category  | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part333E                                 | 09/09/2010           |                    |

## Part 7 of 8

### GENUINE FIRST AID BURN ANTISEPTIC PAIN RELIEF WITH ALOE

benzalkonium chloride, lidocaine cream

## Product Information

|                         |                |
|-------------------------|----------------|
| Item Code (Source)      | NDC:52124-0004 |
| Route of Administration | TOPICAL        |

## Active Ingredient/Active Moiety

| Ingredient Name  | Basis of Strength       | Strength        |
|--|-------------------------|-----------------|
| <b>BENZALKONIUM CHLORIDE</b> (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y) | BENZALKONIUM CHLORIDE   | 0.13 g in 100 g |
| <b>LIDOCAINE HYDROCHLORIDE</b> (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)  | LIDOCAINE HYDROCHLORIDE | 0.5 g in 100 g  |

## Inactive Ingredients

| Ingredient Name                                 | Strength |
|---|----------|
| <b>ALOE VERA LEAF</b> (UNII: ZY81Z83H0X)        |          |
| <b>CETYL ALCOHOL</b> (UNII: 936JST6JCN)         |          |
| <b>DIAZOLIDINYL UREA</b> (UNII: H5RIZ3MPW4)     |          |
| <b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)      |          |
| <b>GLYCERIN</b> (UNII: PDC6A3C0OX)              |          |
| <b>GLYCERYL MONOSTEARATE</b> (UNII: 230OU9XXE4) |          |
| <b>METHYL PARABEN</b> (UNII: A2I8C7HI9T)        |          |
| <b>MINERAL OIL</b> (UNII: T5L8T28FGP)           |          |
| <b>POLYETHYLENE GLYCOLS</b> (UNII: 3WJQ0SDW1A)  |          |
| <b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)      |          |
| <b>PROPYL PARABEN</b> (UNII: Z8IX2SC1OH)        |          |
| <b>WATER</b> (UNII: 059QF0K00R)                 |          |
| <b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)          |          |

## Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|-----------|---------------------|----------------------|--------------------|
|---|-----------|---------------------|----------------------|--------------------|

|   |                  |                   |  |  |
|---|------------------|-------------------|--|--|
| 1 | NDC:52124-0004-1 | 0.9 g in 1 PACKET |  |  |
|---|------------------|-------------------|--|--|

### Marketing Information

| Marketing Category  | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part333A                                 | 04/13/2011           |                    |

### Part 8 of 8

### INSECT STING RELIEF PAD

benzocaine,alcohol swab

### Product Information

|                         |                |
|-------------------------|----------------|
| Item Code (Source)      | NDC:52124-0008 |
| Route of Administration | TOPICAL        |

### Active Ingredient/Active Moiety

| Ingredient Name  | Basis of Strength | Strength         |
|--|-------------------|------------------|
| BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5) | BENZOCAINE        | 0.6 mg in 100 mL |
| ALCOHOL (UNII: 3K9958V90M) (ALCOHOL - UNII:3K9958V90M)       | ALCOHOL           | 60 mL in 100 mL  |

### Inactive Ingredients

| Ingredient Name          | Strength |
|--------------------------|----------|
| WATER (UNII: 059QF0KO0R) |          |

### Packaging

| # | Item Code        | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------|----------------------|--------------------|
| 1 | NDC:52124-0008-1 | 0.5 mL in 1 PACKAGE |                      |                    |

### Marketing Information

| Marketing Category  | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part348                                  | 04/23/2011           |                    |

### Marketing Information

| Marketing Category      | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|-------------------------|--|----------------------|--------------------|
| OTC monograph not final | part333B                                 | 01/09/2013           |                    |



**Labeler** - Genuine First Aid, LLC (619609857)

**Registrant** - GFA Production (Xiamen) Co., Ltd. (421256261)

Revised: 1/2013

Genuine First Aid, LLC