ZICAM COLD AND FLU-LIKE SYMPTOMS ECHINACEA PLUS SAMBUCUS- zinc acetate anhydrous, zinc gluconate, echinacea angustifolia whole, and sambucus nigra flowering top tablet, multilayer CHURCH AND DWIGHT CO, INC.

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

ZICAM® COLD & FLU-LIKE SYMPTOMS Echinacea + Sambucus

Drug Facts

Active Ingredients (in each tablet)	Purpose
Echinacea angustifolia 5x	Helps to relieve cold and flu-like symptoms*
Sambucus nigra (elderberry) 5x	Helps to relieve cold and flu-like symptoms [*]
Zincum aceticum 2x Zincum gluconicum 1x	Reduces duration of the common cold and helps relieve nasal congestion

* Claims based on traditional homeopathic practice, not accepted medical evidence. Not FDA evaluated.

Uses

Reduces duration of the common cold and helps relieve nasal congestion and temporarily relieves

- Fever
- Chills
- Chest Congestion
- Hoarseness
- Cough
- Sore throat
- Body aches

Warnings

If you are allergic or sensitive to zinc, consult a doctor before using.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomitting, consult a doctor promptly.

Stop use and ask a doctor if symptoms, including cough, persist, recur or are accompanied by a fever, rash or persistent headache, or if new symptoms occur. A

persistent cough that lasts for more than one week may be a sign of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- for best results, use at the first sign of cold and flu and continue to use until symptoms completely subside.
- adults and children 12 years of age and older:
 - take 1 tablet at the onset of symptoms.
 - dissolve entire tablet in mouth. Do not chew. Do not swallow whole.
 - repeat every 2 3 hours, not to exceed 7 tablets in 24 hours. Take until symptoms are gone.
 - to avoid minor stomach upset, do not take on an empty stomach.
 - do not eat or drink for 15 minutes after use. Do not eat or drink citrus fruits or juices for 30 minutes before or after use. Otherwise, drink plenty of fluids.
- children under 12 years of age: ask a doctor before use.

Other information

store between 15-29°C (59-84°F)

Inactive ingredients

crospovidone, FD&C Blue No. 2 aluminum lake, FD&C Yellow No. 6 aluminum lake, flavors, magnesium stearate, mannitol, mono-ammonium glycyrrhizinate, silicon dioxide, sodium starch glycolate, stearic acid, sucralose.

Questions? Comments?

call 877-942-2626 toll free

DISTRIBUTED BY CHURCH & DWIGHT CO., INC. EWING, NJ 08628

PRINCIPAL DISPLAY PANEL - 20 Tablet Blister Pack Carton

NEW! ZICAM® COLD & FLU-LIKE SYMPTOMS‡ ECHINACEA + SAMBUCUS HELPS WITH 6

SYMPTOMS‡

✓ COUGH

✓ HOARSENESS

- ✓ SORE THROAT
- ✓ BODY ACHES
- ✓ CHEST CONGESTION
- ✓ FEVER & CHILLS

Plus Zinc to help relieve nasal congestion

HOMEOPATHIC

20 QUICK-DISSOLVE TABLETS

COOLING PEPPERMINT FLAVOR

‡Claims based on traditional homeopathic practice, not accepted medical evidence. Not FDA evaluated.





Zicam® Cold Remedy products are available in a wide range of forms and flavors.

ZICAM®

COLD & FLU-LIKE SYMPTOMS[‡]

ECHINACEA + SAMBUCUS

HELPS WITH

- COUGH
 SORE THROAT
 CHEST CONGESTION
- 🕑 HOARSENESS

SYMPTOMS:

- BODY ACHES
- **G** FEVER & CHILLS

Plus Zinc to help relieve nasal congestion

2

10 MEO PATHIC 20 QUICK-DISSOLVE TABLETS



‡Claims based on traditional homeopathic practice, not accepted medical evidence. Not FDA evaluated

SATISFACTION GUARANTEEL Questions? Comments? call 877-942-2626 toll free www.zicam.com

WWW.ZICAIM.COM Keep Carton For Complete Warnings and information Distributed bychurch & dwight Cd., inc. Ewing. NJ 08528

he first sign of a cold — you have a Pre-Cold®

Take every 2-3 HOURS until symptoms are gone.

SAFETY SEALED. DO NOT USE IF MARKED SEAL ON BLISTER IS DISTURBED OR MISSING. KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION.

This product is not required to go through the FDA's New Drug Application approval process.

Drug Facts

Active Ingredients (in each tablet) Purpose Sambucus nigra (elderberry) 5xHelps to relieve cold and flu-like symptoms‡ Zincum aceticum 2x Reduces duration of the common cold and Zincum gluconicum 1xhelps relieve nasal congestion Uses Reduces duration of the common cold and helps relieve nasal congestion and temporarily relieves -Fever -Chills -Chest Congestion -Hoarseness -Cough -Sore throat -Body aches Warnings If you are allergic or sensitive to zinc, consult a doctor before using. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomitting, consult a doctor promptly. Stop use and ask a doctor if symptoms, including cough, persist, recur or are accompanied by a fever, rash or persistent headache, or if new symptoms occur. A persistent cough that lasts for more than one week may be a sign of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Directions for best results, use at the first sign of cold and flu and continue to use until symptoms completely subside. adults and children 12 years of age and older: take 1 tablet at the onset of symptoms. dissolve entire tablet in mouth. Do not chew. Do not swallow whole. repeat every 2 – 3 hours, not to exceed 7 tablets in 24 hours. Take until symptoms are gone. to avoid minor stomach upset, do not take on an empty stomach. do not eat or drink for 15 minutes after use. Do not eat or drink citrus fruits or juices for 30 minutes before or after use. Otherwise, drink plenty of fluids. children under 12 years of age: ask a doctor before use. Other information store between 15-29°C (59-84°F)

Inactive ingredients crospovidone, FD&C Blue No. 2 aluminum lake, FD&C Yellow No. 6 aluminum lake, flavors, magnesium stearate, mannitol, mono-ammonium glycyrrhizinate, silicon dioxide, sodium starch glycolate, stearic acid, sucralose.

‡Claims based on traditional homeopathic practice, not accepted medical evidence. Not FDA evaluated.

ZICAM COLD AND FLU-LIKE SYMPTOMS ECHINACEA PLUS SAMBUCUS

zinc acetate anhydrous, zinc gluconate, echinacea angustifolia whole, and sambucus nigra flowering top tablet, multilayer

Product Information						
Product Type	HUMAN OTC D	RUG	Item Code (Sou	ırce)	NDC:1023	7-479
Route of Administration	ORAL					
Active Ingredient/Acti	ive Moiety					
Ingredient Name Basis of Strer						Strength
ZINC ACETATE ANHYDROUS (UNII: H2ZEY72PME) (ZINC CATION - UNII:13S1S8SF37) ZINC CATION						2 [hp_X]
ZINC GLUCONATE (UNII: U6W	ZINC GLUCONATE (UNII: U6WSN5SQ1Z) (ZINC CATION - UNII:13S1S8SF37) ZINC GLUCONATE				IATE	1 [hp_X]
ECHINACEA ANGUSTIFOLIA WHOLE (UNII: VB06AV5US8) (ECHINACEA ANGUSTIFOLIA WHOLE - UNII: VB06AV5US8) ECHINACEA ANGUSTIFOLIA WHOLE - UNII: VB06AV5US8)					WHOLE	5 [hp_X]
SAMBUCUS NIGRA FLOWERING TOP (UNII: CT03BSA18U) (SAMBUCUS NIGRA FLOWERING TOP - UNII:CT03BSA18U)SAMBUCUS NIGRA FLOWERING TOP						5 [hp_X]
Inactive Ingredients	Inactive Ingredients					
Ingredient Name Strengt						trength
CROSPOVIDONE (120 .MU.M) (UNII: 68401960MK)						
FD&C BLUE NO. 2ALUMINUM LAKE (UNII: 4AQJ3LG584)						
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)						
MANNITOL (UNII: 30WL53L36A	MANNITOL (UNII: 30WL53L36A)					
AMMONIUM GLYCYRRHIZATE (UNII: 3VRD35U26C)						
MAGNESIUM STEARATE (UNII: 70097M6I30)						
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)						
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)						
STEARIC ACID (UNII: 4ELV7Z 65AP)						
SUCKALUSE (UNII: 96K6UQ32	SUCRALOSE (UNII: 96K6UQ3ZD4)					
Product Characteristi	Product Characteristics					
Color ORA	NGE, BLUE		Score		no score	2

	nape	R	OUND	Size		15mm
Fl	avor	PI	EPPERMINT	Imprint	Code	ZZ
Co	ontains					
P	ackaging					
#					Marketing Start Date	Marketing End Date
1	NDC:10237- 479-20	1 in 1 CA	ARTON		07/01/2023	
1		20 in 1 BLISTER PACK; Type 0: Not a Combination Product				
M	larketing	Infor	mation			
Μ	larketing Marketing Category		mation oplication Number or N Citation	1onograph	Marketing Start Date	Marketing End Date

Labeler - CHURCH AND DWIGHT CO, INC. (001211952)

Establishment						
Name		Address	ID/FEI	Business Operations		
LNK International, Inc.			833150803	MANUFACTURE(10237-479)		
Establishment						
Name	Address	5 ID/FEI		Business Operations		
LNK International, Inc.		967626305	MANUFACTURE	(10237-479), PACK(10237-479)		
Establishment						
Name	Address	s ID/FEI		Business Operations		
LNK International, Inc.		868734088	8 MANUFACTURE	(10237-479) , PACK(10237-479)		
Establishment						
Name	Address	s ID/FEI		Business Operations		
LNK International, Inc.		038154464	MANUFACTURE	(10237-479), PACK(10237-479)		

Establishment						
Name	Address	ID/FEI	Business Operations			
LNK International, Inc.		832867837	MANUFACTURE(10237-479), PACK(10237-479)			

Revised: 6/2023

CHURCH AND DWIGHT CO, INC.