MILK OF MAGNESIA- magnesium hydroxide suspension STRATEGIC SOURCING SERVICES LLC

fos mom

Active ingredient (in each 15 mL tablespoonful)

Magnesium hydroxide 1200 mg

Purpose

Saline laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts over 14 days

Ask a doctor or pharmacist before use if you are

presently taking a prescription drug.

This product may interact with certain prescription drugs.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (800) 222-1222.

Directions

- shake well before use
- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor

• drink a full glass (8 oz) of liquid with each dose

adults and children 12 years and older	2 to 4 tablespoonfuls
children 6 to 11 years	1 to 2 tablespoonfuls
children under 6 years	ask a doctor

Directions for 12 oz

- shake well before use
- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, only use the dosing cup provided
- mL = milliliter

age	dose
adults and children 12 years and over	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

Other information

- each 15 mL tablespoonful contains: magnesium 500 mg
- store at room temperature and avoid freezing
- close cap tightly after use

Inactive ingredients

purified water, sodium hypochlorite

Questions or comments?

Call **833-358-6431**

package Label

NDC 70677-1074-1

COMPARE TO THE ACTIVE INGREDIENT OF PHILLIPS* MILK OF MAGNESIA*

Foster & Thrive™

Milk of Magnesia

Magnesium Hydroxide SALINE LAXATIVE

- · Gentle Overnight Relief · Sugar Free
- Stimulant Free
- Cramp Free



ORIGINAL FLAVOR DOSAGE CUP INCLUDED

12 FL OZ (355 mL)

275-19112-0 **REV 0623**

Drug Facts TAMPER EVIDENT: DO NOT USE IF IMPRINTED NECKBAND IS MISSING OR BROKEN.

Active ingredient (in each 15 mL)

Purpose

Magnesium hydroxide 1200 mg _____

Saline laxative

Use • relieves occasional constipation (irregularity)

. generally produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before use if you have

a magnesium-restricted diet

· kidney disease

· stomach pain, nausea or vomiting

a sudden change in bowel habits that lasts over 14 days

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. This product may interact with certain prescription drugs.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (800) 222-1222.

Directions • shake we before use

- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a ful glass (8 oz) of liquid with each dose
- for accurate dosing, only use the dosing cup provided mL = milliliter

age	dose
adults and children 12 years and over	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

Other information • each 15 mL contains: magnesium 500 mg avoid freezing · store at room temperature tightly closed

Inactive ingredients purified water, sodium hypochlorite

Questions or comments? Call 833-358-6431

*This product is not manufactured or distributed by Bayer Healthcare LLC, owner of the registered trademark Phillips * Milk of Magnesia.

©2023 McKesson Corporation Distributed by: McKesson Corp., via SSSL. Memphis, TN 38141 Money Back Guarantee www.fosterandthrive.com



NDC 70677-1074-2

COMPARE TO THE ACTIVE INGREDIENT OF PHILLIPS** MILK OF MAGNESIA*

Foster & Thrive™

Milk of Magnesia

Magnesium Hydroxide SALINE LAXATIVE

- Gentle Overnight Relief
- Stimulant Free
- Sugar Free
- Cramp Free

ORIGINAL FLAVOR

16 FL OZ (473 mL)

275-19112-0 REV 0623

Drug Facts

TAMPER EVIDENT: DO NOT USE IF IMPRINTED NECKBAND IS MISSING OR BROKEN.

Active ingredient (in each 15 mL tablespoonful)
Magnesium hydroxide 1200 mg

Purpose Saline laxative

Use • relieves occasional constipation (irregularity)

. generally produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before use if you have • kidney disease

a magnesium-restricted diet
 stomach pain, nausea, or vomiting

· a sudden change in bowel habits that lasts over 14 days

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. This product may interact with certain prescription drugs.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product.
 These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

you need to use a raxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a

Poison Control Center right away (800) 222-1222.

Directions • shake well before use

- . do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- . drink a full glass (8 oz) of liquid with each dose

age	dose	
adults and children 12 years and older	2 to 4 tablespoonfuls	
children 6 to 11 years	1 to 2 tablespoonfuls	
children under 6 years	ask a doctor	

Other information • each 15 mL tablespoonful contains: magnesium 500 mg
• store at room temperature and avoid freezing • close cap tightly after use

Inactive ingredients purified water, sodium hypochlorite

Questions or comments? Call 833-358-6431

©2023 McKesson Corporation *This produ Distributed by: McKesson Corp., Bayer via SSSL Memphis, TN 38141 Money Back Guarantee • www.fosterandthrive.com

910-19111-0 Rev 0623

This product is not manufactured or distributed by Bayer Healthcare LLC, owner of the registered trademark Phillips Milk of Magnesia.



MILK OF MAGNESIA

magnesium hydroxide suspension

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:70677-1074

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name

Basis of Strength

MAGNESIUM HYDROXIDE (UNII: NBZ 3QY004S) (MAGNESIUM CATION - UNII: 1200 mg in 15 mL

Inactive Ingredients

Ingredient Name Strength

WATER (UNII: 059QF0KO0R)

SODIUM HYPOCHLORITE (UNII: DY38VHM5OD)

Packaging					
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:70677- 1074-1	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2023	
	2	NDC:70677- 1074-2	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2023	

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M007	06/01/2023		

Labeler - STRATEGIC SOURCING SERVICES LLC (116956644)

Registrant - GCP Laboratories (965480861)

Establishment				
Name	Address	ID/FEI	Business Operations	
GCP Laboratories		965480861	pack(70677-1074) , manufacture(70677-1074)	

Revised: 11/2023 STRATEGIC SOURCING SERVICES LLC