

**VICKS NYQUIL HIGH BLOOD PRESSURE COLD AND FLU- acetaminophen,
dextromethorphan hbr, doxylamine succinate capsule, liquid filled
The Procter & Gamble Manufacturing Company**

VICKS® NyQuil™ HIGH BLOOD PRESSURE COLD & FLU, LiquiCaps™

Drug Facts

Active ingredients (in each liquicap)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Doxylamine succinate 6.25 mg

Active ingredients (in each liquicap) Purpose

Pain reliever/fever reducer
Cough suppressant
Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever
- runny nose & sneezing

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 Liquicaps in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters

- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 8 LiquiCaps per 24 hrs

| | |
|---------------------------------|------------------------------------|
| adults & children 12 yrs & over | 2 LiquiCaps with water every 4 hrs |
| children 4 to under 12 yrs | ask a doctor |
| children under 4 yrs | do not use |

Other information

store at no greater than 25°C (77°F)

Inactive ingredients

D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, lecithin, mica, polyethylene glycol, polyvinyl acetate phthalate, povidone, sorbitol sorbitan solution, titanium dioxide, water

Questions?

1-800-362-1683

TAMPER EVIDENT: DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED.

DIST. BY PROCTER & GAMBLE,

CINCINNATI, OH 45202

PRINCIPAL DISPLAY PANEL - 24 LiquiCaps™

MAX STRENGTH

VICKS®

NyQuil™

HIGH BLOOD PRESSURE

COLD & FLU

HBP

Acetaminophen- Pain reliever/fever reducer

Doxylamine Succinate - Antihistamine

Dextromethorphan HBr - Cough suppressant

DECONGESTANT FREE

24 LIQUICAPS™

| | |
|--|--|
| <p>Drug Facts</p> <p>Active ingredients (in each LiquiCap)</p> <p>Acetaminophen 325 mg Pain reliever/fever reducer</p> <p>Dextromethorphan HBr 10 mg Cough suppressant</p> <p>Doxylamine succinate 6.25 mg Antihistamine</p> <p>Purpose</p> | <p>Drug Facts (continued)</p> <p>Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> • skin reddening • blisters • rash <p>If a skin reaction occurs, stop use and seek medical help right away</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> |
| <p>Uses temporarily relieves common cold/flu symptoms:</p> <ul style="list-style-type: none"> • cough due to minor throat & bronchial irritation • sore throat • headache • minor aches & pains • fever • runny nose & sneezing | <p>Do not use</p> <ul style="list-style-type: none"> • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. |
| <p>Warnings</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take</p> <ul style="list-style-type: none"> • more than 8 Liquicaps in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product | <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> • liver disease • glaucoma • cough that occurs with too much phlegm (mucus) |

| | | | | | | | |
|---|--|---------------------------------|------------------------------------|----------------------------|--------------|----------------------|------------|
| <p>Drug Facts (continued)</p> <ul style="list-style-type: none"> • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema • trouble urinating due to enlarged prostate gland | <p>Drug Facts (continued)</p> <p>reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> | | | | | | |
| <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> • taking sedatives or tranquilizers • taking the blood thinning drug warfarin | <p>Directions</p> <ul style="list-style-type: none"> • take only as directed • do not exceed 8 LiquiCaps per 24 hrs <table border="1"> <tr> <td>adults & children 12 yrs & over</td> <td>2 LiquiCaps with water every 4 hrs</td> </tr> <tr> <td>children 4 to under 12 yrs</td> <td>ask a doctor</td> </tr> <tr> <td>children under 4 yrs</td> <td>do not use</td> </tr> </table> | adults & children 12 yrs & over | 2 LiquiCaps with water every 4 hrs | children 4 to under 12 yrs | ask a doctor | children under 4 yrs | do not use |
| adults & children 12 yrs & over | 2 LiquiCaps with water every 4 hrs | | | | | | |
| children 4 to under 12 yrs | ask a doctor | | | | | | |
| children under 4 yrs | do not use | | | | | | |
| <p>When using this product</p> <ul style="list-style-type: none"> • excitability may occur, especially in children • marked drowsiness may occur • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • alcohol, sedatives, and tranquilizers may increase drowsiness | <p>Other information store at no greater than 25°C (77°F)</p> <p>Inactive Ingredients D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, lecithin, mica, polyethylene glycol, polyvinyl acetate phthalate, povidone, sorbitol sorbitan solution, titanium dioxide, water</p> | | | | | | |
| <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> • pain or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. <p>These could be signs of a serious condition.</p> | <p>Questions? 1-800-362-1683</p> | | | | | | |
| <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of</p> | | | | | | | |

acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:69423-804 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------------------|-----------------|
| DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL) | DOXYLAMINE SUCCINATE | 6.25 mg |
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 10 mg |
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 325 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--|-----------------|
| LECITHIN, SOYBEAN (UNII: 1DI56QDM62) | |
| POLYVINYL ACETATE PHTHALATE (UNII: 58QVG85GW3) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A) | |
| FD&C BLUE NO. 1 (UNII: H3R47K3TBD) | |
| GELATIN (UNII: 2G86QN327L) | |
| GLYCERIN (UNII: PDC6A3C00X) | |
| WATER (UNII: 059QF0K00R) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| POVIDONE (UNII: FZ989GH94E) | |
| D&C YELLOW NO. 10 (UNII: 35SW5USQ3G) | |
| MICA (UNII: V8A1AW0880) | |
| SORBITOL (UNII: 506T60A25R) | |

Product Characteristics

| | | | |
|-----------------|-------|---------------------|----------|
| Color | green | Score | no score |
| Shape | OVAL | Size | 15mm |
| Flavor | | Imprint Code | NQ |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|----------|------------------|---|-----------------------------|---------------------------|
| 1 | NDC:69423-804-24 | 1 in 1 CARTON | 07/11/2023 | |
| 1 | | 24 in 1 BOTTLE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M012 | 07/11/2023 | |

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Establishment

| Name | Address | ID/FEI | Business Operations |
|-----------------------|---------|-----------|------------------------|
| Patheon Softgels Inc. | | 002193829 | manufacture(69423-804) |

Revised: 7/2024

The Procter & Gamble Manufacturing Company